



Europäisches Patentamt
European Patent Office
Office européen des brevets



⑪ Publication number: **0 232 932 B1**

⑫

EUROPEAN PATENT SPECIFICATION

④⑤ Date of publication of patent specification :
02.01.91 Bulletin 91/01

⑤① Int. Cl.⁵: **C07D 253/06, C07D 409/10,
A61K 31/53**

②① Application number : **87200088.0**

②② Date of filing : **21.01.87**

⑤④ **5,6-Dihydro-2-(substituted phenyl)-1,2,4-triazine-3,5(2H,4H)-diones.**

The file contains technical information
submitted after the application was filed and
not included in this specification

③① Priority : **30.01.86 GB 8602342**

④③ Date of publication of application :
19.08.87 Bulletin 87/34

④⑤ Publication of the grant of the patent :
02.01.91 Bulletin 91/01

⑧④ Designated Contracting States :
AT BE CH DE ES FR GB GR IT LI LU NL SE

⑤⑥ References cited :
**EP-A- 0 154 885
EP-A- 0 170 316
FR-A- 2 231 378
US-A- 3 912 723**

⑦③ Proprietor : **JANSSEN PHARMACEUTICA N.V.**
Turnhoutseweg 30
B-2340 Beerse (BE)

⑦② Inventor : **Boeckx, Gustaaf Maria**
Augustijnenstraat 21
B-2360-Oud-Turnhout (BE)
Inventor : **Raeymaekers, Alfons Herman**
Margaretha
Aanbeeldstraat 1
B-2340-Beerse (BE)
Inventor : **Slipido, Victor**
Winterkoningstraat 37
B-2060-Merksem (BE)

EP 0 232 932 B1

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

2-Phenyl-as-triazine-3,5(2H,4H)-diones and their use for controlling coccidiosis have been described in U.S. Patent No. 3,912,723. The phenyl moiety in the said triazines may, inter alia, be substituted with a benzoyl-, an α -hydroxy-phenylmethyl- and a phenylsulfonyl radical.


Substituted 2-phenyl-hexahydro-1,2,4-triazine-3,5-diones and their use for combatting Protozoa have been disclosed in Published Eur. Pat. Application No. 0,154,885.

The 5,6-dihydro-2-phenyl-1,2,4-triazine-3,5(2H,4H)-diones, described in the present application, differ from the hereinabove-mentioned triazinones, by the specific substitution of the 2-phenyl moiety, resulting in 5,6-dihydro-1,2,4-triazine-3,5(2H,4H)-diones which are very effective in destructing or preventing the growth of Protozoa in subjects suffering from such Protozoa.

A number of compounds structurally closely related to the ones the present invention are disclosed in EP-A-0 170 316, some of such compounds being excluded by provisions from the definition of the compounds of formula (II-J) as defined hereafter.

Description of the preferred embodiments :

The present invention is related to 5,6-dihydro-2-(substitutedphenyl)-1,2,4-triazine-3,5(2H,4H)-diones having the formula



the pharmaceutically acceptable acid addition, metal or amine substitution salts, and stereochemically isomeric forms thereof, wherein :

Ar is thienyl, halo substituted thienyl, naphthalenyl or a radical of formula



R is hydrogen, C_{1-6} alkyl, cyclo C_{3-6} alkyl, aryl or (aryl) C_{1-6} alkyl ;

R^1 is cyano or a radical of formula $-C(=X)-Y-R^2$;

said X being O or S,

Y being O, S, NR^{10} or a direct bond ; R^2 being hydrogen, aryl, C_{3-6} cycloalkyl or C_{1-6} alkyl optionally substituted with aryl, hydroxy, amino, mono- and di(C_{1-6} alkyl)amino, piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C_{1-6} alkyl)-piperazinyl, 4-(C_{1-6} alkyl-carbonyl)-piperazinyl, 4-(C_{1-6} alkyloxycarbonyl)-piperazinyl or 4-((aryl) C_{1-6} alkyl)-piperazinyl ; and where Y is a direct bond, R^9 may also be halo ; R^{10} is hydrogen, C_{1-6} alkyl or (aryl) C_{1-6} alkyl ; or R^9 and R^{10} taken together with the nitrogen atom bearing said R^9 and R^{10} may form a piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C_{1-6} alkyl)piperazinyl, 4-(C_{1-6} alkylcarbonyl)-piperazinyl, 4-(C_{1-6} alkyloxycarbonyl)-piperazinyl or a 4-((aryl) C_{1-6} alkyl)-piperazinyl radical ;

R^2 , R^3 , R^6 , R^7 and R^8 are each independently hydrogen, halo, trifluoromethyl, C_{1-6} alkyl, hydroxy, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyloxy, mercapto, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, C_{1-6} alkylsulfinyl, (trifluoromethyl)-sulfonyl, cyano, nitro, amino, mono- and di(C_{1-6} alkyl)amino, or (C_{1-6} alkylcarbonyl)amino ;

R^4 and R^5 are each independently hydrogen, aryl, cyclo C_{3-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, (aryl) C_{2-6} alkenyl or C_{1-6} alkyl optionally substituted with aryl, hydroxy, amino, mono- and di(C_{1-6})amino, piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C_{1-6} alkyl)-piperazinyl, 4-(C_{1-6} alkylcarbonyl)-piperazinyl, 4-(C_{1-6} alkyloxy-carbonyl)-piperazinyl or 4-((aryl) C_{1-6} alkyl)-piperazinyl ;

and R^5 may also be C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl, (aryl) C_{1-6} alkyloxycarbonyl or (aryl)carbonyl ; wherein aryl is phenyl, optionally substituted with up to 3 substituents each independently selected from

the group consisting of halo, C₁₋₆ alkyl, C₁₋₆ alkyloxy, trifluoromethyl, hydroxy, mercapto, C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, trifluoromethylsulfonyl, cyano, nitro, amino, mono- and di (C₁₋₆alkyl)amino and (C₁₋₆ alkylcarbonyl)amino.

In the foregoing definitions the term "halo" is generic to fluoro, chloro, bromo and iodo; "C₁₋₆ alkyl" is meant to include straight and branched saturated hydrocarbon radicals, having from 1 to 6 carbon atoms, such as, for example, methyl, ethyl, 1-methylethyl, 1,1-dimethylethyl, propyl, butyl, pentyl and hexyl "cyclo C₃₋₆ alkyl" embraces cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; "C₂₋₆ alkenyl" is meant to include straight and branch chained hydrocarbon radicals containing one double bond and having from 2 to 6 carbon atoms such as, for example, ethenyl, 3-propenyl and 2-butenyl "C₂₋₆ alkynyl" is meant to include straight and branch chained hydrocarbon radicals containing one triple bond and having from 2 to 6 carbon atoms such as, for example, ethynyl, 3-propynyl and 2-butylnyl


Preferred compounds within the invention are those wherein Ar is halothienyl or a radical of formula (a) wherein R⁶ and R⁷ are, each independently, hydrogen, halo, trifluoromethyl, C₁₋₆ alkyloxy, hydroxy or C₁₋₆ alkyl; R⁸ is hydrogen; R is hydrogen, C₁₋₆ alkyl, phenyl or halophenyl; R² and R³ are, each independently, hydrogen, halo, trifluoromethyl or C₁₋₆ alkyl; and R⁴ is hydrogen or C₁₋₆ alkyl.

Particularly preferred compounds within the invention are those preferred compounds wherein Ar is a radical of formula (a) wherein R⁶ is halo, R⁷ and R⁸ are hydrogen, R is hydrogen or C₁₋₆ alkyl, R² and R³ independently are halo or hydrogen.

More particularly preferred compounds within the invention are those particularly preferred compounds wherein R⁶ is 4-chloro, R is hydrogen, R² is 2-chloro, R³ is 6-chloro or hydrogen and R⁴ is hydrogen.

The most preferred compounds within the invention are 2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile and 2-chloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl) benzeneacetonitrile and the pharmaceutically acceptable acid addition, metal or amine substitution salts thereof.

The compounds of formula (I) may conveniently be prepared by a reduction reaction of the corresponding 1,2,4-triazine-3,5-(2H,4H)-dione of formula (II), or an acid-addition salt, metal or amine substitution salt form thereof, thus preparing a compound of formula (I) wherein R⁵ is hydrogen, said compounds being represented by the formula (I-a), and if desired, subsequently reacting the compounds of formula (I-a) with a reagent R^{5-a}-W (III), thus preparing compounds of formula (I), wherein R⁵ is other than hydrogen, said compounds being represented by the formula (I-b). In (III) W represents an appropriate reactive leaving group such as, for example, halo, e.g., chloro, bromo or iodo, or a sulfonyloxy group, e.g. methylsulfonyloxy or 4-methylphenyl-sulfonyloxy and R^{5-a} has the previously defined meaning of R⁵, provided that it is not hydrogen.




The said reduction reaction is conveniently conducted following art-known procedures for converting a 1,2,4-triazine-3,5-(2H,4H)-dione into a 5,6-dihydro-1,2,4-triazine-3,5-(2H,4H)-dione moiety. A number of such procedures are described in for example the Published Eur. Pat. Application No. 0,154,885 and the references cited therein.


Said reduction reaction may for example be conducted by contacting the starting material of formula (II) with hydrogen in the presence of an appropriate catalyst such as, for example, Raney-nickel, platinum, palladium, platinum(IV) oxide, and the like. Preferably, said reduction reaction is conducted by reacting the starting material (II) with zinc in acetic acid or tin(II) chloride in hydrochloric acid, optionally in the presence of a reaction-inert organic solvent or mixture of such solvents such as, for example, a lower alkanol, e.g. methanol or ethanol; a hydrocarbon, e.g. methylbenzene or dimethylbenzene; a ketone, e.g. 2-propanone, 1-butanone; an ether, e.g. tetrahydrofuran, 1,2-dimethoxyethane, 1,4-dioxane, an ester, e.g. ethyl acetate; N,N-dimethylformamide, N,N-dimethylacetamide; pyridine; acetic acid. Higher temperatures may be used to enhance the reaction rate.

The alkylation reaction of (I-a) with the reagent R^5-W may be conducted following art-known N-alkylation procedures. The alkylation reaction is conveniently conducted in an inert organic solvent such as, for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene and dimethylbenzene, a lower alkanol, e.g., methanol, ethanol and 1-butanol a ketone, e.g., 2-propanone and 4-methyl-2-pentanone an ether, e.g., 1,4-dioxane, 1,1'-oxybisethane and tetrahydrofuran N,N-dimethylformamide (DMF); N,N-dimethylacetamide (DMA); nitrobenzene; dimethyl sulfoxide (DMSO) and 1-methyl-2-pyrrolidinone. The addition of an appropriate base such as, for example, an alkali metal carbonate or hydrogen carbonate, sodium hydride or an organic base such as, for example N,N-diethylethanamine or N-(1-methylethyl)-2-propanamine may be utilized to pick up the acid which is liberated during the course of the reaction. In some circumstances the addition of an iodide salt, preferably an alkali metal iodide, is appropriate. Somewhat elevated temperatures may enhance the rate of the reaction.

The compounds of formula (I) wherein R^1 is cyano, said compounds being represented by the formula




may alternatively be prepared by converting the hydroxy function of a triazinedione of formula



into a nitrile function.

The conversion of (IV) into (I-c) may be effected by art-known procedures. For example, by first converting the hydroxy function into a suitable leaving group and subsequently converting the said leaving group in the thus obtained intermediate having the formula




into a nitrile function.

In (V) W has the meaning of an appropriate reactive leaving group such as, for example, halo, e.g., chloro, bromo or iodo, or a sulfonyloxy group, e.g. methylsulfonyloxy or 4-methylphenylsulfonyloxy.

For example, where W represents chloro, the intermediates (V) may be prepared by reacting (IV) with thionyl chloride in a suitable reaction-inert solvent.

The conversion of (V) into (I-c) may be effected, e.g., by reacting (V) with a cyanide, such as, for example, an alkalimetal cyanide, e.g. potassium cyanide, sodium cyanide; copper cyanide; silver cyanide and the like, if desired, in the presence of an appropriate solvent.


The compounds of formula (I) can also be prepared by reacting a 5,6-dihydro-1,2,4-triazine-3,5(2H,4H)-dione having the formula (VII) with a reagent of formula (VI) following art-known procedures for arylating an amine functionality.



In (VI) W represents an appropriate reactive leaving group as defined hereinabove and preferably is halo. The reaction of (VI) with (VII) is most conveniently conducted in an appropriate reaction-inert solvent, preferably at higher temperature and in the presence of a suitable base. Suitable reaction inert solvents may, for example, be aromatic hydrocarbons, e.g., benzene, methylbenzene and dimethylbenzene; halogenated hydrocarbons, e.g. trichloromethane and trichloroethane; dipolar aprotic solvents such as N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide and the like.


The compounds of formula (I) can also be converted into each other following art-known procedures of functional group transformation. Some examples will be cited hereinafter.

In order to simplify the structural representations of the compounds of formula (I) in the schemes illustrating these group transformation procedures, the




group will hereafter be represented by the symbol D.

For example, the compounds of formula (I) wherein R¹ is cyano, said compounds being represented by the formula (I-c) may partially or completely be hydrolysed, thus yielding compounds of formula (I) wherein the radical R¹ is a carboxyl or an aminocarbonyl group, the former compounds being represented by the formula (I-e), the latter by (I-d). The said partial hydrolysis reaction is preferably conducted in an aqueous acidic medium, e.g. an aqueous sulfuric, hydrochloric or phosphoric acid solution, at room temperature or at slightly increased temperature. Complete hydrolysis is accomplished when increasing either the reaction temperature or the reaction time or both. In the said complete hydrolysis reaction it may be advantageous to add a second acid to the reaction mixture, e.g. acetic acid. In turn, the compounds of formula (I-d) can further be hydrolysed to obtain compounds of formula (I-e) by treating the starting compounds of formula (I-d) with an aqueous acidic solution in the same way as for obtaining (I-e) from (I-c).



20 Furthermore, the compounds of formula (I-c) can be converted to the corresponding compounds of formula (I) wherein R^1 is an aminothioxomethyl group, said compounds being represented by the formula (I-f), by reacting (I-c) with hydrogen sulfide, preferably in a suitable solvent such as pyridine, optionally in the presence of an organic base such as a trialkylamine, e.g. triethylamine.




35 The acids of formula (I-e) can be converted to the corresponding acylhalides of formula (I-g) by treatment with a suitable halogenating agent such as, for example, thionyl chloride, pentachlorophosphorane, sulfur chloride. The thus obtained acylhalides of formula (I-g) can further be derivatized to the corresponding amides of formula (I-h) by reacting the starting acylhalide compounds of formula (I-g) with a suitable amine optionally in the presence of an appropriate solvent such as an ether, e.g. tetrahydrofuran, acetonitrile, trichloromethane or dichloromethane.

40 Or, the said acylhalides of formula (I-g) can conveniently be converted to the corresponding aryl or alkyl ketones of formula (I-i), respectively (I-j), by reacting (I-g) with benzene or substituted benzene in the presence of a Lewis acid catalyst such as aluminum chloride, respectively with a metal alkyl, e.g. methyl lithium, butyl lithium, optionally in the presence of a suitable catalyst, e.g. copper (I) iodide, or with a complex metal alkyl, in a suitable solvent, e.g. tetrahydrofuran.

45 The acylhalides of formula (I-g) can further be converted to the corresponding esters of formula (I-k) by a suitable alcoholysis reaction. Or, the acids of formula (I-f) can be esterified following art-known procedures, e.g. by treating the starting acids with an alkylhalide in a suitable solvent in the presence of a base, e.g. in N,N-dimethylformamide or N,N-dimethylacetamide in the presence of an alkalimetal carbonate, or by reacting (I-g) with the appropriate alcohol in the presence of a reagent capable of forming an ester functionality e.g. N,N'-methanetetraylbis[cyclohexanamine].

50

55




25 The compounds of formula (I), wherein either R⁴ or R⁵ is hydrogen may be converted to the compounds of formula (I) having an R⁴ and/or R⁵ other than hydrogen by N-alkylating or N-acylating the starting compounds following art-known procedures, e.g. following the same procedures as described hereinabove for the preparation of (I-b) starting from (I-a).

30 The compounds of formula (I) which contain an Ar moiety which is a phenyl radical substituted with one or more alkoxy radicals may be converted to the corresponding alkylcarbonyloxy compounds by treating the starting compounds with an alkanolic acid in the presence of anhydrous hydrohalic acid, e.g. hydrobromic acid in acetic acid. The said alkylcarbonyloxy compounds can in turn be converted to the corresponding hydroxyphenyl compounds by a suitable hydrolysis reaction, e.g. by treatment with an aqueous hydrohalic solution.


35 A number of intermediates and starting materials in the foregoing preparations are known compounds which may be prepared according to art-known methodologies of preparing said or similar compounds and others are new. A number of such preparation methods will be described hereinafter in more detail.

Several of the intermediates of formula (II) are known compounds and their synthesis is described in the Published Eur. Pat. Application No.0,170,316.

40 They may generally be prepared by cyclizing an intermediate of formula




and eliminating the group E of the thus obtained dione



In the intermediates (VIII) L has the meaning of an appropriate leaving group such as C₁₋₆ alkyloxy, halo and the like. The group E, as described in the intermediate (VIII) and the triazinedione (IX), represents an appropriate electron attracting group which may conveniently be eliminated from the dione (IX) such as, for example, a carboxyl, a sulfonyloxy, a sulfinyloxy group or a precursor and/or derivative thereof, e.g. an ester, an amide, a cyanide, a C₁₋₆ alkylsulfonyloxy, phenylsulfonyloxy, C₁₋₆ alkylphenylsulfonyloxy and halophenylsulfonyloxy and the like groups.


A particularly suitable process for preparing intermediates of formula (II) consists of cyclizing an intermediate of formula (VIII-a) and eliminating the E¹ functionality in the thus obtained intermediate of formula (IX-a). In (VIII-a) and (IX-a) E¹ represents a cyano, C₁₋₆ alkyloxycarbonyl or amido group.



The cyclization reaction may be effected following art-known cyclization procedures as described, for example, in Monatshefte der Chemie, 94, 258-262 (1963), e.g. by heating the starting compound of formula (VIII-a) over its melting point, or by refluxing a mixture of (VIII-a) with a suitable solvent such as, for example, an aromatic hydrocarbon, e.g. benzene, methylbenzene, or dimethylbenzene, an acid, e.g. acetic acid, optionally in the presence of base, e.g. potassium acetate, sodium acetate and the like.


The elimination of the E¹ functionality may be effected following art-known procedures as described, for example, in Monatshefte der Chemie, 96, 134-137 (1965), e.g. by converting (IX-a) into a carboxylic acid (X) in a suitable acidic reaction medium such as acetic acid, aqueous hydrochloric acid solutions or mixtures thereof. Elevated temperatures may enhance the rate of the reaction.

The thus obtained carboxylic acids of formula



10 may be converted into an intermediate of formula (II) by art-known decarboxylation reaction procedures, e.g. by heating the carboxylic acid (X) or by heating a solution of (X) in 2-mercaptoacetic acid as described, for example, in US Patent No. 3,896,124.


15 The intermediates of formula (VIII) may generally be prepared by reacting a diazonium salt of formula (XI) with a reagent of formula (XII).



X⁻, as described in (XI) has the meaning of an appropriate anion and E and L, as described in (XII), have the previously defined meanings.

30 The reaction of (XI) with (XII) may conveniently be conducted in a suitable reaction medium as described, for example, in Monatshefte der Chemie, 94, 694-697 (1963). Suitable reaction mediums are, for example, aqueous sodium acetate solutions, pyridine and the like.


35 The starting diazonium salts (XI) may be derived from a corresponding amine of formula (XIII) following art-known procedures, e.g. by reacting the latter with an alkalimetal or earth alkaline metal nitrite, e.g. sodium nitrite, in a suitable reaction medium.



In the hereinabove-described reaction scheme Mⁿ⁺ is a alkalimetal or earth alkaline metal kation and n is the integer 1 or 2.


50 The amines of formula (XIII) may be prepared following procedures analogous to those described in U.S. Patent No. 4,005,218.

55 The triazin-4(1H)-ones of formula (IV) may conveniently be prepared by reducing the corresponding 1,2,4-triazine-3,5 (2H,4H)-diones of formula (XIV) following the same procedures as described hereinabove for the preparation of (I) starting from (II) and, if desired, by further N-alkylating the thus obtained triazin-4(1H)-one of formula (IV-a) with a reagent R⁵-W (III) following the same procedures described hereinabove for the preparation of (I-b) starting from (I-a).




The triazinediones of formula (XIV) may be prepared following the procedures described in US Patent No. 3,912,723.

The intermediates of formula (II) wherein R¹ is cyano, said intermediates being represented by the formula




may alternatively be prepared by converting the hydroxy function of a triazinedione of formula



into a nitrile function.

The conversion of (XV) into (II-a) may be effected by art-known procedures. For example, by first converting the hydroxy function into a suitable leaving group and subsequently converting the said leaving group in the thus obtained intermediate having the formula



10

into a nitrile function.

In (XVI) W has the previously defined meaning of an appropriate reactive leaving group.


For example, where W represents chloro, the intermediates (XVI) may be prepared by reacting (XV) with thionyl chloride in a suitable reaction-inert solvent.

15

The conversion of (XVI) into (II-a) may be effected, e.g., by reacting (XVI) with a cyanide, such as, for example, an alkali metal cyanide, e.g. potassium cyanide, sodium cyanide and copper cyanide and silver cyanide, if desired, in the presence of an appropriate solvent.

The intermediates of formula (II) may alternatively be prepared by reacting a 1,2,4-triazine-3,5-(2H,4H)-dione of formula (XVII) with an aromatic compound of formula (VI) following the same procedures as described hereinabove for the preparation of (I) starting from (VI) with (VII).

20



The intermediates of formula (II) can also be converted into each other following art-known procedures of functional group transformation. A number of such group transformations are represented by the following scheme. In order to simplify the structural representations of the intermediates of formula (II) in these schemes, the



35

40

45

50


55



25

The conversion of (II-a) to (II-b), of (II-a) to (II-c) and of (II-b) to (II-c) can conveniently be done following the same procedures as described hereinabove for the conversion of (I-c) to (I-d), (I-c) to (I-e) and of (I-d) to (I-e).

A further series of such grouptransformations can be represented by the following scheme.



The conversion of (II-a) to (II-d), of (II-d) to (II-e), of (II-e) to (II-f), of (II-d) to (II-i), of (II-e) to (II-i), of (II-e) to (II-h) and of (II-e) to (II-g) can conveniently be done following the same procedures as described hereinabove for the conversion of (I-c) to (I-f), of (I-f) to (I-g), of (I-g) to (I-h), of (I-f) to (I-k), of (I-g) to (I-k), of (I-g) to (I-j) and of (I-g) to (I-i).


The intermediates of formula (II), wherein R^4 is hydrogen may be converted to the compounds of formula (II) having an R^4 other than hydrogen by N-alkylating the starting compounds following art-known procedures, e.g. the procedures described hereinabove for the preparation of (I-b) starting from (I-a).

The intermediates of formula (II) which contain a phenyl radical substituted with one or more alkyloxy radicals may be converted to the corresponding alkylcarbonyloxy compounds which in turn may be converted to the corresponding hydroxyphenyl compounds following the same procedures as described hereinabove for the analogous conversions of the compounds of formula (I).

An additional feature of the present invention comprises the fact that a number of intermediates mentioned in the foregoing preparation schemes are novel compounds.

More particularly, the intermediates of formula (II) wherein R^1 is other than cyano, and the intermediates of formula (II) wherein R^1 is cyano and in the latter case one of the following conditions is met :


- a) R is other than hydrogen, C_{1-6} alkyl, cyclo C_{3-6} alkyl or aryl ; or
- b) Ar is other than a radical of formula (a) wherein in said radical of formula (a) R^6 is other than hydrogen, halo, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio or C_{1-6} alkylsulfonyl ; or
- c) R^2 is other than hydrogen, halo, trifluoromethyl or C_{1-6} alkyl ; or
- d) R^4 is other than hydrogen, said intermediates being represented by the formula



and the pharmaceutically acceptable acid-addition, metal or amine substitution salts, and stereochemically isomeric forms thereof are novel compounds.

Preferred, particularly preferred and more particularly preferred intermediates of formula (II-j) are those of which the preferred, particularly preferred and more particularly preferred compounds of formula (I) can be derived, i.e. those intermediates wherein Ar , r , r^1 , r^2 , r^3 and r^4 are as defined hereinabove for the corresponding radicals Ar , R , R^1 , R^2 , R^3 and R^4 of the preferred, particularly preferred and more particularly preferred compounds of formula (I).

Furthermore, the pharmaceutically acceptable metal or amine substitution salts of those intermediates of formula (II), wherein Ar is a radical of formula (a), and wherein R^6 , R^7 and R^8 are each independently hydrogen, halo, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio, or C_{1-6} alkylsulfonyl ; R^2 and R^3 are each independently hydrogen, halo, trifluoromethyl or C_{1-6} alkyl ; and wherein R^1 is cyano, R^4 is hydrogen and wherein R is hydrogen, C_{1-6} alkyl, cyclo C_{3-6} alkyl or phenyl optionally substituted with up to 3 substituents each independently selected from the group consisting of halo, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio and C_{1-6} alkylsulfonyloxy, said compounds being represented by the formula



and the stereochemically isomeric forms thereof are also novel compounds.

Preferred are those metal or amine salts of those intermediates of formula (II-k) wherein R^{6-a} is halo and more preferably is chloro, R^{7-a} and R^{8-a} are hydrogen, R^a is hydrogen or C_{1-6} alkyl and more preferably is hydrogen and R^{2-a} and R^{3-a} independently are hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkyloxy and more pref-

erably R^{2a} is chloro and R^{3a} is hydrogen or chloro.

Both the intermediates of formula (II-j) and (II-k) are not only novel compounds useful in the preparation of the compounds of formula (I), but they also possess anti-protozoal and more particularly anti-coccidial activity.

5 The compounds of formula (I) and the intermediates of formula (II) may be converted to their therapeutically active non-toxic acid addition salt forms by treatment with appropriate acids, such as, for example, inorganic acids, such as hydrohalic acid, e.g. hydrochloric, hydrobromic and sulfuric acid, nitric acid and phosphoric acid or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxy-propanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, 10 ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic and 4-amino-2-hydroxybenzoic. Conversely the salt form can be converted by treatment with alkali into the free base form.

15 The compounds of formula (I) and the intermediates of formula (II), containing one or more acidic protons, may also be converted to their therapeutically active non-toxic metal or amine substitution salt forms by treatment with appropriate organic or inorganic bases. Appropriate inorganic bases may, for example, be ammonia or bases derived from alkali or earth alkaline metals, e.g. alkali metal or earth alkaline metal oxides or hydroxides such as lithium hydroxide, sodium hydroxide, potassium hydroxide, magnesium hydroxide, calcium hydroxide and calcium oxide; alkalimetal or earth alkaline metal hydrides, e.g. sodium hydride and potassium hydride; alkalimetal hydrogen carbonates or carbonates, e.g. sodium carbonate, 20 potassium carbonate, sodium hydrogen carbonate and calcium carbonate. Appropriate organic bases may, for example be primary, secondary and tertiary aliphatic and aromatic amines such as, for example, methylamine, ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine, pyrrolidine, piperidine, morpholine, N-methylmorpholine, trimethylamine, tripropylamine, quinuclidine, pyridine, quinoline, isoquinoline, diethanolamine and 1,4-diazabicyclo[2,2,2]octane; or quaternary ammonium bases e.g. tetramethylammonium hydroxide, trimethylbenzylammonium hydroxide, triethylbenzylammonium hydroxide, tetraethylammonium hydroxide, and trimethylethylammonium hydroxide.

25 It is obvious from formulae (I) and (II) that the compounds of the present invention have an asymmetric carbon atom. Consequently, these compounds may exist under two different enantiomeric forms. Pure enantiomeric forms of the compounds of formula (I) and the intermediates of formula (II) may be obtained by the application of art-known procedures.

30 The compounds of formula (I) and the intermediates of formula (II), the pharmaceutically acceptable acid addition salts, metal or amine substitution salts and the possible stereochemically isomeric forms thereof are useful agents in combatting Protozoa. For example, said compounds are found to be active against a wide variety of said Protozoa such as, for example, Sarcodina, Mastigophora, Ciliophora and Sporozoa.

35 The compounds of formula (I) and the intermediates of formula (II), the pharmaceutically acceptable acid addition salts, metal or amine substitution salts and the possible stereochemically isomeric forms thereof are especially useful agents in combatting Rhizopoda such as, for example, Entamoeba; and Mastigophora such as, for example, Trichomonas, e.g. Trichomonas vaginalis, Histomonas, e.g. Histomonas maleagridis, and Trypanosoma spp.

40 In view of their potent activity in combatting Protozoa the compounds of this invention constitute useful tools for the destruction or prevention of the growth of Protozoa and more particularly they can effectively be used in the treatment of subjects suffering from such Protozoa.

45 In view of the potent activity in combatting Protozoa this invention provides valuable compositions comprising the compounds of formula (I) and the intermediates of formula (II), the acid addition salts, metal or amine substitution salts or possible stereochemically isomeric forms thereof, as the active ingredient in a solvent or a solid, semi-solid or liquid diluent or carrier, and, in addition, it provides an effective method of combatting Protozoa by use of an effective anti-protozoal amount of such compounds of formula (I) and the 50 intermediates of formula (II), or acid addition salts or metal or amine substitution salts thereof. Anti-protozoal compositions comprising an effective amount of an active compound of formula (I) or an active intermediate of formula (II), either alone or in combination with other active therapeutic ingredients, in admixture with suitable carriers may be readily prepared according to conventional pharmaceutical techniques for the usual 55 routes of administration.

Preferred compositions are in dosage unit form, comprising per dosage unit an effective quantity of the active ingredient in admixture with suitable carriers. Although the amount of the active ingredient per unit dosage may vary within rather wide limits, dosage units comprising from about 10 to about 2000 mg of the

active ingredient are preferred.

In view of the anti-protozoal properties of the compounds of formula (I) and the intermediates of formula (II) it is evident that the present invention provides a method of inhibiting and/or eliminating the development of Protozoa in warm-blooded animals suffering from diseases caused by one or more of these Protozoa by the administration of an antiprotozoal effective amount of a compound of formula (I) and intermediates of formula (II), a pharmaceutically acceptable acid addition salt, metal or amine substitution salts or a possible stereochemically isomeric form thereof.

More particularly, in view of their extremely potent activity in combatting Coccidia the compounds of this invention are very useful in the destruction or prevention of the growth of Coccidia in warm-blooded animals. Consequently, the compounds of formula (I) and the intermediates of formula (II), the acid addition salts, metal or amine substitution salts and possible stereochemically isomeric forms thereof are particularly useful anti-coccidial agents as well as coccidiostatics.

Due to their useful anti-coccidial and coccidiostatic activity the subject compounds may be administered in combination with any solid, semi-solid or liquid diluent or carrier as described hereinabove. Additionally, due to their useful coccidiostatic activity the subject compounds may be mixed with any kind of feed supplied to warm-blooded animals although it may also be administered while dissolved or suspended in the drinking water.

The following examples are intended to illustrate the present invention. Unless otherwise stated all parts therein are by weight.

EXAMPLES

A) Preparation of Intermediates

Example 1

Procedure A

To a stirred mixture of 16 parts of 2-[3-chloro-4-[(2,4-dichlorophenyl)hydroxymethyl]phenyl]-1,2,4-triazine-3,5(2H,4H)-dione and 150 parts of trichloromethane are added dropwise, during a period of 5 minutes, 16 parts of thionyl chloride. Upon completion, stirring is continued for 3 hours at reflux temperature. The reaction mixture is evaporated in vacuo. Methylbenzene is added and the whole is evaporated again, yielding 2-[3-chloro-4-[chloro(2,4-dichlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2H,4H)-dione.

A mixture of 12 parts of 2-[3-chloro-4-[chloro(2,4-dichlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2H,4H)-dione and 5.4 parts of copper cyanide is stirred and heated first for 3 hours at 130°C and for 3 hours at 180°C. After cooling, the precipitated product is dissolved in a mixture of trichloromethane and methanol (90:10 by volume). The inorganic precipitate is filtered off and the filtrate is evaporated in vacuo. The residue is purified by column chromatography over silica gel using first a mixture of trichloromethane and acetonitrile (93:7 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is dissolved in 20 parts of N,N-dimethyl-formamide and 25 parts of 1,1'-oxybisethane. The product is allowed to crystallize, filtered off and dried, yielding 2-chloro- α -(2,4-dichlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile (Intermediate 1).

Procedure B

To a stirred mixture of 68 parts of 1-chloro-2-methoxy-4-nitrobenzene, 230 parts of a sodium hydroxide solution 50%, 5 parts of N,N,N-triethylbenzenemethanaminium chloride and 360 parts of tetrahydrofuran is added dropwise, during a 5 minutes period, a solution of 43.2 parts of 4-chlorobenzeneacetonitrile in 90 parts of tetrahydrofuran. Upon completion, stirring is continued for 4 hours at 60°C. The reaction mixture is poured into 2000 parts of crushed ice. The whole is acidified with concentrated hydrochloric acid. The layers are separated. The aqueous phase is extracted with dichloromethane. The combined extracts are dried, filtered and evaporated. The residue is crystallized from 2,2'-oxybispropane. The product is filtered off and dried, yielding α -(4-chlorophenyl)-2-methoxy-4-nitrobenzeneacetonitrile.

A mixture of 8.1 parts of α -(4-chlorophenyl)-2-methoxy-4-nitrobenzeneacetonitrile, 2 parts of a solution of thiophene in methanol 4% and 200 parts of methanol is hydrogenated at normal pressure and at 50°C with 2 parts of platinum-on-charcoal catalyst 5%. After the calculated amount of hydrogen is taken up, the catalyst is filtered off and the filtrate is evaporated in vacuo. The residue is stirred in a small amount of 2-propanol. The product is filtered off and dried, yielding 4-amino- α -(4-chlorophenyl)-2-methoxyben-

zeneacetonitrile.


To a stirred and cooled (5-10°C) mixture of 5.6 parts of 4-amino- α -(4-chlorophenyl)-2-methoxybenzeneacetonitrile, 6.2 parts of concentrated hydrochloric acid and 50 parts of acetic acid is added dropwise, during a 15 minutes period, a solution of 1.25 parts of sodium nitrite in 10 parts of water at about 10°C. Upon completion, the whole is stirred for 60 minutes and then 3.6 parts of anhydrous sodium acetate and 2.8 parts of ethyl (2-cyanoacetyl)carbamate are added and stirring is continued for 2 hours at room temperature. The reaction mixture is poured into 250 parts of water. The product is filtered off, washed with water and dissolved in a mixture of trichloromethane and methanol (90 :10 by volume). The organic layer is dried, filtered and evaporated. The residue is purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95 :5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated in vacuo, yielding ethyl [2-[[4-[(4-chlorophenyl)cyanomethyl]-3-methoxyphenyl]hydrazono]-2-cyanoacetyl]carbamate.

A mixture of 8.3 parts of ethyl [2-[[4-[(4-chlorophenyl)cyanomethyl]-3-methoxyphenyl]hydrazono]-2-cyanoacetyl]carbamate, 1.77 parts of anhydrous potassium acetate and 100 parts of acetic acid is stirred for 2 hours at reflux temperature. The reaction mixture is evaporated in vacuo. The residue is stirred in water. The product is filtered off and dissolved in trichloromethane. The organic layer is dried, filtered and evaporated. The residue is purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90 :10 by volume) as eluent. The pure fractions are collected and the eluent is evaporated in vacuo, yielding 2-[4-[(4-chlorophenyl)cyanomethyl]-3-methoxyphenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carbonitrile.


A mixture of 4 parts of 2-[4-[(4-chlorophenyl)cyanomethyl]-3-methoxyphenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carbonitrile, 24 parts of concentrated hydrochloric acid and 40 parts of acetic acid is stirred and refluxed for 3 hours. The reaction mixture is evaporated in vacuo and the residue is stirred in water. The product is extracted with a mixture of trichloromethane and methanol (90 :10 by volume). The extract is dried, filtered and evaporated, yielding 2-[4-[(4-chlorophenyl)cyanomethyl]-3-methoxyphenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid.

A mixture of 4.2 parts of 2-[4-[(4-chlorophenyl)cyanomethyl]-3-methoxyphenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid and 13 parts of 2-mercaptoacetic acid is stirred and heated for 2 hours at 175°C. After cooling, 150 parts of water are added. The aqueous phase is decanted and the remaining oil is stirred in water. The whole is treated with sodium hydrogen carbonate. The product is extracted with a mixture of trichloromethane and methanol (90 :10 by volume). The extract is dried, filtered and evaporated. The residue is purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97 :3 by volume) as eluent. The pure fractions are collected and the eluent is evaporated in vacuo. The residue is crystallized from 8 parts of acetonitrile. The product is filtered off, washed with 2,2'-oxybispropane and dried, yielding α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-3-methoxybenzeneacetonitrile (intermediate 2).

Following the same procedures there are further prepared :



int. no.	R ⁶	R ⁷	R ⁸	R ²	R ³
3	2-Cl	4-Cl	6-Cl	2-Cl	H
4	4-Br	H	H	2-Cl	H
5	4-Br	H	H	2-Cl	6-Cl
6	4-CH ₃ COO-	H	H	2-Cl	6-Cl
7	4-OH	H	H	2-Cl	6-Cl
8	4-Cl	H	H	2-OH	H
9	4-CH ₃ S	H	H	2-Cl	H
10	4-CH ₃ S	H	H	2-Cl	6-Cl
11	4-CH ₃ S	H	H	2-CH ₃	6-CH ₃
12	4-CH ₃ S	3-CH ₃	H	2-Cl	H
13	4-CH ₃ S	3-CH ₃	H	2-Cl	6-Cl
14	4-CH ₃ SO	H	H	2-Cl	H
15	4-CH ₃ SO	H	H	2-Cl	6-Cl
16	4-CH ₃ SO ₂	H	H	2-Cl	H
17	4-CH ₃ SO ₂	H	H	2-Cl	6-Cl
18	4-HS	H	H	2-Cl	H
19	4-HS	H	H	2-Cl	6-Cl



Comp.	Ar	R	R ²	R ³
20	2-Cl-5-thienyl	H	Cl	Cl
21	2-Cl-5-thienyl	H	Cl	H
22	2-Cl-5-thienyl	CH ₃	Cl	Cl
23	2-Cl-5-thienyl	CH ₃	Cl	H
24	1-naphthalenyl	H	Cl	Cl
25	1-naphthalenyl	H	Cl	H

Example 2

To 30 parts of a sulfuric acid solution in water (90 : 10 by volume) were added portionwise during a period of 5 minutes 2 parts of 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile at room temperature. Upon completion, stirring was continued for 2 hours at 80°C. The reaction mixture was poured into ice water. The product was filtered off, washed with water and purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95 : 5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was stirred in 2,2'-oxybispropane. The product was filtered off and dried, yielding 1.1 parts (54%) of 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetamide ; mp. 160.7°C (Intermediate 26).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared :
2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetamide ; mp. 276.4°C (Intermediate 27).

Example 3

To a stirred mixture of 9.2 parts of concentrated sulfuric acid, 5 parts of acetic acid and 5 parts of water were added 1.5 parts of 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile at room temperature. The whole was stirred and refluxed for 18 hours. The reaction mixture was poured into 100 parts of ice water. The product was filtered off, washed with water and purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and acetic acid (95 : 4 : 1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was stirred in 2,2'-oxybispropane. The product was filtered off and dried, yielding 0.9 parts (59%) of 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetic acid ; mp. 196.3°C (Intermediate 28).

Example 4

A mixture of 13.2 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetamide, 648 parts of concentrated hydrochloric acid and 200 parts of acetic acid was stirred and refluxed for 224 hours. The resulting product was filtered off, washed with water and taken up in 100 parts of water. After treatment with a sodium hydroxide solution, the resulting solution was acidified with concentrated hydrochloric acid. The product was filtered off and purified by column chromatography over

silica gel using a mixture of methylbenzene, tetrahydrofuran and acetic acid (70 :30 :1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 3.8 parts (27.8%) of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetic acid ; mp. 219.5°C (intermediate 29).

Example 5

A mixture of 6 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile, 1.5 parts of N,N-diethylethanamine and 40 parts of pyridine was stirred at room temperature. Gaseous hydrogen sulfide was bubbled through the mixture during 24 hours. The solvent was evaporated *in vacuo* and the residue was stirred in water. The precipitated product was filtered off, stirred in 2-propanol and filtered off again. The product was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97 :3 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was crystallized from 16 parts of acetonitrile. The product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 1.4 parts (21.1%) of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneethanethioamide ; mp. 262.7°C (intermediate 30).

Following the same procedure there is prepared :
2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneethanethioamide (intermediate 31).

Example 6

A mixture of 2 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetic acid and 48 parts of thionyl chloride was stirred for 1 hour at reflux temperature. The reaction mixture was evaporated, yielding 2.28 parts (100%) of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetyl chloride as a residue (intermediate 32).

Following the same procedure there is prepared :
2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetyl chloride (intermediate 33).

Example 7

A mixture of 2.28 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetylchloride and 18 parts of piperidine was stirred for 17 hours at room temperature (exothermic reaction). After the addition of water, the solution was acidified with hydrochloric acid. The product was extracted with a mixture of trichloromethane and methanol (95 :5 by volume). The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and acetic acid (95 :4 :1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was washed with 2,2'-oxybispropane and dried, yielding 1 part (44.0%) of 1-[2-(4-chlorophenyl)-2-[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]acetyl]piperidine ; mp. 216.9°C (intermediate 34).

Example 8

A mixture of 2.28 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetylchloride, 4.5 parts of pyrrolidine and 40 parts of acetonitrile was stirred for 17 hours at room temperature. After evaporation *in vacuo*, the residue was taken up in water and the mixture was acidified with hydrochloric acid. The product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95 :5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was dried *in vacuo* for 48 hours at 110°C, yielding 0.8 parts (36.2%) of 1-[2-(4-chlorophenyl)-2-[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]-acetyl]pyrrolidine ; mp. 153.9°C (intermediate 35).

Example 9

To a stirred mixture of 10 parts of 1-methylpiperazine in 45 parts of tetrahydrofuran was added dropwise a solution of 6.7 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetyl chloride in 45 parts of tetrahydrofuran during a period of 5 minutes. Upon complete addition, stirring was continued for 2 hours at room temperature. After evaporation *in vacuo*, the residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90 :10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was boiled in acetonitrile. After cooling, the precipitated product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 4.8 parts (62.8%) of 1-[2-(4-chlorophenyl)-2-[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]acetyl]-4-methylpiperazine; mp. 261.5°C (intermediate 36).

Following the same procedure there was also prepared :
2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-N-methylbenzeneacetamide; mp. 278.7°C (intermediate 37).

Following the same procedure there are further prepared :
2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-N-methylbenzeneacetamide (intermediate 38).
4-acetyl-1-[2-(4-chlorophenyl)-2-[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]acetyl]piperazine (intermediate 39).
1-[2-(4-chlorophenyl)-2-[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]acetyl]-4-(phenylmethyl)piperazine (intermediate 40).

Example 10

A mixture of 4.7 parts of aluminium trichloride and 67.5 parts of benzene was stirred in an ice bath till a temperature of $\pm 10^\circ\text{C}$. A solution of 4.9 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetyl chloride in 22.5 parts of benzene was added dropwise during a period of 15 minutes at this low temperature (exothermic reaction). Upon complete addition, stirring was continued for 20 hours at room temperature. The reaction mixture was poured into 500 parts of ice water and the whole was acidified with concentrated hydrochloric acid. The product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by filtration over silica gel using a mixture of trichloromethane and methanol (95 :5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was further purified three times by column chromatography over silica gel using first a mixture of trichloromethane and methanol (97 :3 by volume) and then a mixture of trichloromethane and methanol (99 :1 by volume) and finally a mixture of trichloromethane and ethyl acetate (92.5 :7.5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was crystallized from 8 parts of ethanol. The product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 0.7 parts (13.0%) of 2-[3,5-dichloro-4-[1-(4-chlorophenyl)-2-oxo-2-phenylethyl]phenyl]-1,2,4-triazine-3,5(2H,4H)-dione; mp. 143.0°C (intermediate 41).

Example 11

To a stirred and cooled (-70°C , 2-propanone/ CO_2 bath) solution of 5.7 parts of copper(I) iodide in 67.5 parts of tetrahydrofuran were added dropwise 37.5 parts of a methyl lithium solution 1.6 M in 1,1'-oxybisethane during a period of 15 minutes under nitrogen atmosphere. Upon complete addition, stirring was continued for 30 minutes at this low temperature. A mixture of 4.45 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetyl chloride and 22.5 parts of tetrahydrofuran was added dropwise during a period of 30 minutes at -65°C . Upon completion, stirring was continued first for 2 hours at -60°C and then for 1 hour at -20°C . A saturated ammonium chloride solution in water was added dropwise (exothermic reaction). The precipitate was filtered off and from the filtrate, the organic layer was dried, filtered and evaporated. The residue was purified three times by column chromatography over silica gel : twice using a mixture of trichloromethane and methanol (95 :5 and 98 :2 by volume) and then a mixture of trichloromethane and ethyl acetate (92.5 :7.5 by volume) as eluents. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was stirred in acetonitrile. The product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 0.8 parts (18.8%) of 2-[3,5-dichloro-4-[1-(4-chlorophenyl)-2-oxopropyl]phenyl]-1,2,4-triazine-3,5(2H,4H)-dione; mp. 208.4°C (intermediate 42).

Example 12

A mixture of 5.53 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetyl chloride and 160 parts of methanol was stirred for 1 hour at reflux temperature. After evaporation, water was added to the residue and the product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and acetic acid (95 : 4 : 1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was further purified by column chromatography over silica gel using a mixture of trichloromethane, hexane and methanol (45 : 45 : 10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 0.9 parts (17.6%) of methyl 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetate ; mp. 121.1°C (intermediate 43).

Following the same procedure there is also prepared : methyl 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetate (intermediate 44).

Example 13


To a stirred mixture of 8.5 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetic acid, 5.5 parts of potassium carbonate and 45 parts of N,N-dimethylformamide were added 8.52 parts of iodomethane at room temperature. After stirring for 2 hours at 40°C, the reaction mixture was evaporated *in vacuo*. The residue was stirred in water. The precipitated product was filtered off and dissolved in trichloromethane (the remaining water was separated). The organic layer was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (99 : 1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was crystallized from acetonitrile. The product was filtered off (the filtrate was set aside) and dried, yielding a first fraction of 1.9 parts (20.9%) of methyl 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-4-methyl-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetate. The filtrate, which was set aside (see above), was evaporated *in vacuo*, yielding a second fraction of 4 parts (44%) of methyl 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-4-methyl-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetate as a residue. Total yield : 5.9 parts (64.9%) of methyl 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-4-methyl-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetate ; mp. 173.4°C (intermediate 45).

Example 14

To a stirred mixture of 4 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-benzeneacetonitrile, 1.4 parts of potassium carbonate and 22.5 parts of N,N-dimethylformamide were added 2.84 parts of iodomethane at room temperature. The reaction mixture was stirred for 1.5 hours at 40°C. After evaporation *in vacuo*, the residue was taken up in water. The precipitated product was filtered off and washed with water. After crystallization from acetonitrile, the product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 2.5 parts (59.2%) of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-4-methyl-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile ; mp. 159.7°C (intermediate 46).

Following the same procedure there were also prepared : 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-4-(phenylmethyl)-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile ; mp. 128.0°C (intermediate 47) ; 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-4-methyl-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetamide (intermediate 48) ; and (E)-2,6-dichloro- α -(4-chlorophenyl)-4-[4,5-dihydro-3,5-dioxo-4-(3-phenyl-2-propenyl)-2H-1,2,4-triazin-2-yl]benzeneacetonitrile ; mp. 159.2°C (intermediate 49).

Following the same procedures there are further prepared :



Int. no.	R ⁴
50	-C ₂ H ₅
51	-C ₃ H ₇ -1
52	-C ₄ H ₉ -n
53	-CH ₂ -CH=CH
54	-CH ₂ -C≡CH
55	-CH ₂ CH ₂ -C ₆ H ₅

Example 15

A mixture of 13 parts of 2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-α-(4-methoxyphenyl)benzeneacetonitrile and 300 parts of acetic acid, saturated with hydrogen bromide was stirred for 24 hours at 90°C. The reaction mixture was poured into 500 parts of ice water. The precipitated product was filtered off, washed with water and dissolved in trichloromethane. The remaining aqueous layer was removed. The organic layer was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and ethyl acetate (80:20 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was purified twice by column chromatography over silica gel using a mixture of trichloromethane and ethyl acetate (85:15 by volume) as eluent. The first fraction was collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 2 parts (14.4%) of 4-[cyano[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-phenyl]-methyl]phenol acetate (ester); mp. 221.5°C (intermediate 56).

A mixture of 2.3 parts of 4-[cyano[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]-methyl]phenol acetate(ester) and 50 parts of a hydrochloric acid solution 4 N was stirred for 4 hours at reflux temperature. The precipitate was filtered off (the filtrate was set aside), washed successively with water, 2-propanol and 2,2'-oxybispropane and dried. The precipitate was combined with the filtrate, which was set aside (see above) and the solvent was evaporated. The residue was stirred and refluxed for 4 hours. After evaporated *in vacuo*, the residue was dissolved in a mixture of trichloromethane and methanol (90:10 by volume). The solution was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated *in vacuo*. The residue was dissolved in a mixture of acetonitrile and 2,2'-oxybispropane (5:20 by volume). The crystallized product was filtered off and dried, yielding 1 part (48.4%) of 2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-α-(4-hydroxyphenyl)benzeneacetonitrile; mp. 209.9°C (intermediate 57).

Example 16

To a stirred mixture of 5 parts of 2,6-dichloro-α-(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-

2(3H)yl)benzeneacetonitrile in 40 parts of water were added 5 parts of a solution of 9.6 parts of sodium hydroxide in 100 parts of water under nitrogen atmosphere. The whole was stirred for 10 minutes. The precipitate was filtered off. The product in the filtrate was allowed to crystallize. The product was filtered off, washed with water and dried over weekend at 50°C, yielding 2.4 parts (44.6%) of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile, sodium salt monohydrate ; mp. 213.1°C (intermediate 58).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared :

2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile, potassium salt sesquihydrate ; mp. 150.7°C (intermediate 59).

B) Preparation of Final Compounds.

Example 17


To a stirred and refluxed solution of 1.5 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-benzeneacetic acid in 100 parts of acetic acid were added 3 parts of zinc. Stirring was continued for 30 minutes at reflux temperature. The zinc salts were filtered off and the filtrate was evaporated. The residue was washed with water whereupon the solid product was filtered off and dissolved in a mixture of trichloromethane and methanol (90 :10 by volume). The solution was dried, filtered and evaporated. The residue was washed with ethyl acetate and 2,2'-oxybispropane and dried, yielding 0.98 parts (67.2%) of 2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetic acid ; mp. 194.2°C (compound 1).

Example 18

To a stirred and refluxed mixture of 1.5 parts of 2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)- α -(4-fluorophenyl)- α -methylbenzeneacetonitrile and 75 parts of acetic acid were added portionwise 3 parts of zinc during a period of 30 minutes. Upon complete addition, stirring was continued for 3 hours at reflux. The reaction mixture was filtered while hot and the filtrate was concentrated in vacuo to 10 parts of its volume. Water was added to the concentrate. The precipitated product was filtered off and purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95 :5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated in vacuo. The residue was crystallized from acetonitrile. The product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 0.8 parts (53.6%) of 2-chloro- α -(4-fluorophenyl)- α -methyl-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile ; mp 122.5°C (compound 2).


Example 19

Following the procedures of examples 17 and 18 using the appropriate starting materials the following compounds are prepared :




comp. no.	R ¹	R ⁷	R ⁸	R	R ²	R ³	m. p. °C
3	H	H	H	CH ₃	H	H	-
4	4-Cl	H	H	CH ₃	2-Cl	H	-
5	4-Cl	H	H	CH ₃	2-CF ₃	H	-
6	4-Cl	H	H	4-Cl-C ₆ H ₄	2-Cl	H	-
7	4-Cl	H	H	C ₃ H ₇ -n	2-Cl	H	-
8	4-Cl	H	H	C ₄ H ₉ -n	2-Cl	H	-
9	3-CF ₃	4-Cl	H	CH ₃	2-Cl	H	-
10	4-Cl	H	H	CH ₃	2-Cl	6-Cl	-
11	4-Cl	H	H	H	2-Cl	H	168.0
12	4-Cl	H	H	CH ₃	2-Cl	6-CH ₃	211.1
13	4-Cl	H	H	CH ₃	2-Cl	5-CH ₃	-
14	4-Cl	H	H	H	2-Cl	6-Cl	231.2
15	4-Cl	H	H	H	2-Cl	6-CH ₃	-
16	4-F	H	H	H	2-Cl	H	-
17	4-CH ₃	H	H	H	2-Cl	H	-
18	4-F	H	H	CH ₃	2-F	H	165.6
19	4-F	H	H	H	2-Cl	6-Cl	-
20	4-F	H	H	H	2-Cl	6-CH ₃	-
21	4-F	H	H	H	2-CH ₃	6-CH ₃	-
22	4-Cl	H	H	H	2-CH ₃	6-CH ₃	-
23	H	H	H	H	2-Cl	6-Cl	-
24	4-CH ₃ O	H	H	H	2-Cl	6-Cl	201.9
25	2-Cl	4-Cl	H	H	2-Cl	H	-
26	2-Cl	4-Cl	6-Cl	H	2-Cl	H	-

comp. no.	R ¹	R ⁷	R ⁸	R	R ²	R ³	m. p. °C
27	4-Cl	H	H	H	2-CH ₃ O	H	-
28	4-Br	H	H	H	2-Cl	H	-
29	4-Br	H	H	H	2-Cl	6-Cl	-
30	4-CH ₃ COO	H	H	H	2-Cl	6-Cl	-
31	4-OH	H	H	H	2-Cl	6-Cl	-
32	4-Cl	H	H	H	2-OH	H	-
33	4-CH ₃ S	H	H	H	2-Cl	H	-
34	4-CH ₃ S	H	H	H	2-Cl	6-Cl	-
35	4-CH ₃ S	H	H	H	2-CH ₃	6-CH ₃	-
36	4-CH ₃ S	3-CH ₃	H	H	2-Cl	H	-
37	4-CH ₃ S	3-CH ₃	H	H	2-Cl	6-Cl	-
38	4-CH ₃ SO ₂	H	H	H	2-Cl	6-Cl	-
39	4-CH ₃ SO ₂	H	H	H	2-Cl	H	-
40	4-CH ₃ ^O S-	H	H	H	2-Cl	H	-
41	4-CH ₃ ^O S-	H	H	H	2-Cl	6-Cl	-
42	4-HS	H	H	H	2-Cl	6-Cl	-
43	4- ^H S	H	H	H	2-Cl	H	-



Comp.	R ¹	R ³	R ⁴	m.p. °C
44	CONH-CH ₃	6-Cl	H	227.8
45	CONH ₂	6-Cl	H	247.7
46	CO-N(CH ₂) ₄ -N-CH ₃	6-Cl	H	192.1
47	COOH	H	H	-
48	CONH ₂	H	H	-
49	CONH-CH ₃	H	H	-
50	COOCH ₃	H	H	-
51	COOCH ₃	6-Cl	H	-
52	CSNH ₂	6-Cl	H	-
53	CSNH ₂	H	H	-
54	CO-N(CH ₂) ₆ -N-CH ₃	6-Cl	H	-
55	CO-N(CH ₂) ₄ -N-CH ₃	6-Cl	H	-
56	CO-N(CH ₂) ₄ -N-CH ₃	6-Cl	H	-
57	CO-N(CH ₂) ₄ -N-CO-CH ₃	6-Cl	H	-
58	CO-N(CH ₂) ₄ -N-CH ₂ C ₆ H ₅	6-Cl	H	-
59	CO-C ₆ H ₅	6-Cl	H	202.5
60	CO-CH ₃	6-Cl	H	-
61	CN	6-Cl	CH ₃	oil
62	CN	6-Cl	C ₂ H ₅	-
63	CN	6-Cl	C ₃ H ₇ -1	-
64	CN	6-Cl	C ₄ H ₉ -n	-
65	COOCH ₃	6-Cl	CH ₃	123.0
66	CO-NH ₂	6-Cl	CH ₃	220.9

Comp.	R ¹	R ³	R ⁴	m.p. °C
67	CN	6-Cl	-CH ₂ -CH=CH	-
68	CN	6-Cl	-CH ₂ -C≡CH	-
69	CN	6-Cl	-CH ₂ -CH=CH-C ₆ H ₅ (E-form)	175.4
70	CN	6-Cl	-CH ₂ -C ₆ H ₅	186.4
71	CN	6-Cl	-CH ₂ CH ₂ -C ₆ H ₅	



Comp.	Ar	R	R ²	R ³
72	2-Cl-5-thienyl	H	Cl	Cl
73	2-Cl-5-thienyl	H	Cl	H
74	2-Cl-5-thienyl	CH ₃	Cl	Cl
75	2-Cl-5-thienyl	CH ₃	Cl	H
76	1-naphthalenyl	H	Cl	Cl
77	1-naphthalenyl	H	Cl	H

Example 20

A mixture of 4 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-4-methyl-3,5-dioxo-1,2,4-triazin-2-(1H-yl))benzeneacetonitrile, 12 parts of acetic acid anhydride and 36 parts of methylbenzene was stirred for 90 hours at reflux temperature. After cooling, 16 parts of methanol were added. The solvent was evaporated in vacuo. The residue was stirred in water and the product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using trichloromethane as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from ethanol. The precipitated product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 2.0 parts (42.9%) of 1-acetyl-2-[3,5-dichloro-4-[(4-chlorophenyl)cyanomethyl]phenyl]-1,6-dihydro-4-methyl-1,2,4-triazine-3,5(2H,4H)-dione; mp. 178.1°C (compound 78).


Example 21

To a stirred mixture of 3 parts of 2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-4-methyl-3,5-dioxo-1,2,4-triazin-2(1H-yl))-benzeneacetonitrile and 20 parts of pyridine were added dropwise 1.7 parts of benzoyl chloride during a period of 5 minutes at room temperature and under nitrogen atmosphere. Upon complete addition, stirring was continued overnight at room temperature. The reaction mixture was evaporated in vacuo and the residue was stirred in water. The product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified twice by column chromatography over

silica gel using trichloromethane as eluent. The pure fractions were collected and the eluent was evaporated in vacuo. The residue was crystallized from 8 parts of acetonitrile. The product was filtered off, washed with 2,2'-oxybispropane and dried, yielding 0.8 parts (20.2%) of 1-benzoyl-2[3,5-dichloro-4-[(4-chlorophenyl)-cyanomethyl]phenyl]-1,6-dihydro-4-methyl-1,2,4-triazine-3,5(2H,4H)-dione ; mp. 148.4°C (compound 79).

Example 22

Following the same procedures of examples 20 and 21 there are further prepared :



Comp.	R ³	R ⁴	R ⁵
80	Cl	CH ₃	CH ₃
81	Cl	CH ₂ C ₆ H ₅	CH ₂ C ₆ H ₅
82	H	CH ₃	CO-C ₆ H ₅
83	H	CH ₃	CO-CH ₃
84	Cl	CH ₃	CH ₂ C ₆ H ₅
85	Cl	CH ₃	C ₂ H ₅

Example 23

To a stirred mixture of 5 parts of 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetonitrile in 40 parts of water are added 5 parts of a solution of 9.6 parts of sodium hydroxide in 100 parts of water under nitrogen atmosphere. The whole is stirred for 10 minutes. The precipitate is filtered off. The product in the filtrate is allowed to crystallize. The product is filtered off, washed with water and dried for 48 hours at 50°C, yielding 2-chloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile, sodium salt ; (compound 86).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared :

2-chloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile, potassium salt (compound 87) ;

2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile, sodium salt (compound 88) ;

2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile, potassium salt (compound 89) ;

2,6-dichloro- α -(4-chlorophenyl)-4-(3,4,5,6-tetrahydro-3,5-dioxo-1,2,4-triazin-2(1H)-yl)benzeneacetonitrile, ammonium salt (compound 90) ;

C) Pharmacological Examples

The strong anti-protozoal activity of the compounds of formula (I) and the intermediates of formula (II), the pharmaceutically acceptable acid addition salts, metal or amine substitution salts and the possible stereochemically isomeric forms thereof is clearly evidenced by the data obtained in the following experiments, which data are only given to illustrate the useful anti-protozoal properties of all the compounds embraced within the invention and not to limit the invention either with respect to the scope of susceptible Protozoa nor with respect to the scope of formula (I) or of formula (II).

Example 13**Outline of anticoccidial efficacy test against Eimeria tenella.**

- 5 Hisex chickens were fed with a commercial basal ration not containing a coccidiostatic agent.
 Eighteen-day-old chickens were sorted in groups of two birds. Water was supplied automatically and medicated feed was supplied ad libitum from the day of infection (day 0) until the seventh day (not included) after infection. Unmedicated feed was supplied ad libitum to two groups of four birds for uninfected and infected controls.
- 10 Unmedicated feed was a commercial basal ration not containing a coccidiostatic agent. Medicated feed was prepared from unmedicated feed by thoroughly mixing the latter with an amount of the tested compound.
- On day 0 the birds were inoculated orally with 10^5 sporulated oocysts of Eimeria tenella. On day 5 the faecal score was determined and graded :
- 15 0 = no blood spots
 1 = one or two blood spots
 2 = three to five blood spots
 3 = more than five blood spots
- On the seventh day oocysts production is determined by collecting the feces and oocyst count per gram feces (OPG) and the birds are weighed.
- 20 In table 1 the first column shows the average relative weight gain in percent compared with the non-infected controls. The second column shows the average faecal score and the third column illustrates the average oocyst count.

25

30

35

40

45


50

55

int./comp. No.	dose of tested compound in ppm in feed	average relative weight gain	average faecal score	average oocyst count (OPG) x 1000
int. 27	100	107	0	0
int. 29	100	98	0	0
	10	93	0.5	0
int. 30	10	104	0	0
int. 42	10	102	0	0
int. 43	100	96	0	0
	10	98	0	0
int. 46	10	96	0	0
	1	98	0.5	0
int. 58	0.5	101	0	0
	0.1	100	0	1
int. 59	100	98	0	0
	1	96	0	0
	0.5	97	0	0
comp. 1	10	101	1	0
comp. 2	10	100	0	0
comp. 12	10	95	0	0
comp. 14	10	100	0	0
	1	100	0	0
	0.1	101	0	0
comp. 18	10	95	0.5	0
comp. 24	10	92	0	0

Claims


1. A chemical compound having the formula



a pharmaceutically acceptable acid-addition, metal or amine substitution salt, or a stereochemically iso-

meric form thereof, wherein :

Ar is thienyl, halo substituted thienyl, naphthalenyl or a radical of formula



R is hydrogen, C₁₋₆ alkyl, cyclo C₃₋₆ alkyl, aryl or (aryl)C₁₋₆ alkyl ;

R¹ is cyano or a radical of formula -C(=X)-Y-R⁹ ;

15 said X being O or S,

Y being O, S, NR¹⁰ or a direct bond ; R⁹ being hydrogen, aryl, C₃₋₆ cycloalkyl or C₁₋₆ alkyl optionally substituted with aryl, hydroxy, amino, mono- and di(C₁₋₆ alkyl)amino, piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C₁₋₆ alkyl)-piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ alkoxycarbonyl)-piperazinyl or 4-((aryl) C₁₋₆ alkyl)-piperazinyl ; and where Y is a direct bond, R⁹ may also be halo ; R¹⁰ is hydrogen, C₁₋₆ alkyl or (aryl) ; C₁₋₆ alkyl ; or R⁹ and R¹⁰ taken together with the nitrogen atom bearing said R⁹ and R¹⁰ may form a piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C₁₋₆ alkyl)piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ alkoxycarbonyl)-piperazinyl or 4-((aryl) C₁₋₆ alkyl)-piperazinyl radical ;

20 R², R³, R⁶, R⁷ and R⁸ are each independently hydrogen, halo, trifluoromethyl, C₁₋₆ alkyl, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkylcarbonyloxy, mercapto, C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, (trifluoromethyl)-sulfonyl, cyano, nitro, amino, mono- and di(C₁₋₆ alkyl)amino, or (C₁₋₆ alkylcarbonyl)amino ;

25 R⁴ and R⁵ are each independently hydrogen, aryl, cyclo C₃₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, (aryl)C₂₋₆ alkenyl or C₁₋₆ alkyl optionally substituted with aryl, hydroxy, amino, mono- and di(C₁₋₆ alkyl)amino, piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C₁₋₆ alkyl)-piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl or 4-((aryl) C₁₋₆ alkyl)-piperazinyl ;

30 and R⁵ may also be C₁₋₆ alkylcarbonyl, C₁₋₆ alkylcarbonyl,


(aryl)C₁₋₆ alkylcarbonyl or (aryl)carbonyl ; wherein aryl is phenyl, optionally substituted with up to three substituents each independently selected from the group consisting of halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, trifluoromethyl, hydroxy, mercapto, C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, trifluoromethylsulfonyl, cyano, nitro, amino, mono- and di(C₁₋₆ alkyl)amino and (C₁₋₆ alkylcarbonyl)amino.

35 2. A chemical compound according to claim 1 wherein Ar is halothienyl or a radical of formula (a) wherein R⁶ and R⁷ are, each independently, hydrogen, halo, trifluoromethyl, C₁₋₆ alkoxy, hydroxy or C₁₋₆ alkyl ; R⁸ is hydrogen ; R is hydrogen, C₁₋₆ alkyl, phenyl or halophenyl ; R² and R³ are, each independently, hydrogen, halo, trifluoromethyl or C₁₋₆ alkyl ; and R⁴ is hydrogen or C₁₋₆ alkyl.

3. A chemical compound according to claim 2 wherein Ar is a radical of formula (a) wherein R⁶ is halo, R⁷ and R⁸ are hydrogen, R is hydrogen or C₁₋₆ alkyl, R² and R³ independently are halo or hydrogen.


4. A chemical compound according to claim 3 wherein R⁶ is 4-chloro, R is hydrogen, R² is 2-chloro, R³ is 6-chloro or hydrogen and R⁴ is hydrogen.

5. A chemical compound having the formula



55 a pharmaceutically acceptable acid-addition, metal or amine substitution salt, or a stereochemically isomeric form thereof, wherein :

Ar' is thienyl, halo substituted thienyl, naphthalenyl or a radical of formula



5

10

r is hydrogen, C₁₋₆ alkyl, cyclo C₃₋₆ alkyl, aryl or (aryl)C₁₋₆ alkyl ;

r¹ is cyano or a radical of formula -C(=X)-Y-R⁹ ;

said X being O or S,

Y being O, S, NR¹⁰ or a direct bond ; R⁹ being hydrogen, aryl, C₃₋₆ cycloalkyl or C₁₋₆ alkyl optionally substituted with aryl, hydroxy, amino, mono- and di(C₁₋₆ alkyl)amino, piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C₁₋₆ alkyl)-piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ alkyloxy carbonyl)-piperazinyl or 4-((aryl) C₁₋₆ alkyl)-piperazinyl ; and where Y is a direct bond, R⁹ may also be halo ; R¹⁰ is hydrogen, C₁₋₆ alkyl or (aryl) C₁₋₆ alkyl ; or R⁹ and R¹⁰ taken together with the nitrogen atom bearing said R⁹ and R¹⁰ may form a piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C₁₋₆ alkyl)piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ alkyloxycarbonyl)-piperazinyl or a 4-((aryl) C₁₋₆ alkyl)-piperazinyl radical ;

20

r², r³, r⁶, r⁷ and r⁸ are each independently hydrogen, halo, trifluoromethyl, C₁₋₆ alkyl, hydroxy, C₁₋₆ alkyloxy, C₁₋₆ alkylcarbonyloxy, mercapto, C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, (trifluoromethyl)-sulfonyl, cyano, nitro, amino, mono- and di (C₁₋₆ alkyl) amino, or (C₁₋₆ alkylcarbonyl)-amino ;

r⁴ is hydrogen, aryl, cyclo C₃₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, (aryl)C₂₋₆ alkenyl or C₁₋₆ alkyl optionally substituted with aryl, hydroxy, amino, mono- and di (C₁₋₆ alkyl) amino, piperidinyl, pyrrolidinyl, 4-morpholinyl, piperazinyl, 4-(C₁₋₆ alkyl)-piperazinyl, 4-(C₁₋₆ alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ alkyloxy-carbonyl)-piperazinyl or 4-((aryl) C₁₋₆ alkyl)-piperazinyl ;

25

wherein aryl is phenyl, optionally substituted with up to three substituents each independently selected from the group consisting of halo, C₁₋₆ alkyl, C₁₋₆ alkyloxy, trifluoromethyl, hydroxy, mercapto, C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, trifluoromethylsulfonyl, cyano, nitro, amino, mono- and di (C₁₋₆ alkyl) amino and (C₁₋₆ alkylcarbonyl) amino ;

30

provided that, when r¹ is cyano then one of the following conditions is met :

a) r is other than hydrogen, C₁₋₆ alkyl, cyclo C₃₋₆ alkyl or aryl ; or

b) Ar' is other than a radical of formula (a') wherein in said radical of formula (a') r⁶ is other than hydrogen, halo, trifluoromethyl, C₁₋₆ alkyl, C₁₋₆ alkyloxy, C₁₋₆ alkylthio or C₁₋₆ alkylsulfonyl ; or

35

c) r² is other than hydrogen, halo, trifluoromethyl or C₁₋₆ alkyl ; or

d) r⁴ is other than hydrogen.

6. A chemical compound according to claim 5 wherein Ar' is halothienyl or a radical of formula (a') wherein r⁶ and r⁷ are, each independently, hydrogen, halo, trifluoromethyl, C₁₋₆ alkyloxy, hydroxy or C₁₋₆ alkyl ; r⁸ is hydrogen ; r is hydrogen, C₁₋₆ alkyl, phenyl or halophenyl ; r² and r³ are, each independently, hydrogen, halo, trifluoromethyl or C₁₋₆ alkyl ; and r⁴ is hydrogen or C₁₋₆ alkyl.


40

7. A chemical compound according to claim 6 wherein Ar' is a radical of formula (a') wherein r⁶ is halo, r⁷ and r⁸ are hydrogen, r is hydrogen or C₁₋₆ alkyl, r² and r³ independently are halo or hydrogen.

8. A chemical compound according to claim 7 wherein r⁶ is 4-chloro, r is hydrogen, r² is 2-chloro, r³ is 6-chloro or hydrogen and r⁴ is hydrogen.

45

9. A pharmaceutically acceptable metal or amine substitutional salt of a chemical compound having the formula



50

55

or a stereochemically isomeric form thereof, wherein R^{6a} , R^{7a} and R^{8a} are each independently hydrogen, halo, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkylthio or C_{1-6} alkylsulfonyl; R^{2a} and R^{3a} are each independently hydrogen, halo, trifluoromethyl or C_{1-6} alkyl; and R^a is hydrogen, C_{1-6} alkyl, cyclo C_{3-6} alkyl or phenyl optionally substituted with up to 3 substituents each independently selected from the group consisting of halo, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkylthio and C_{1-6} alkylsulfonyloxy.

10. A pharmaceutically acceptable metal or amine substitution salt according to claim 9, wherein R^{6a} is halo, R^{7a} and R^{8a} are hydrogen, R^a is hydrogen or C_{1-6} alkyl and R^{2a} and R^{3a} independently are hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy.

11. A pharmaceutically acceptable metal or amine substitution salt according to claim 10 wherein R^{6a} is chloro, R^a is hydrogen, R^{2a} is chloro and R^{3a} is hydrogen or chloro.

12. A pharmaceutical composition comprising a suitable pharmaceutical carrier and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 11.

13. An anti-protozoal composition comprising an inert carrier and as active ingredient an effective anti-protozoal amount of a compound as claimed in any one of claims 1 to 11.

14. A method of preparing a pharmaceutical composition, characterized in that a therapeutically effective amount of a compound as defined in any of claims 1 to 11 is intimately mixed with suitable pharmaceutical carriers.


15. A compound as claimed in any one of claims 1 to 11 for use as a medicine.

16. A compound as claimed in any one of claims 1 to 11 for use as an anti-protozoal medicine.


17. A process for preparing a chemical compound as claimed in any one of claims 1 to 4, characterized

by

a) reducing an intermediate of formula




thus obtaining a compound of formula




optionally followed by a N-alkylation reaction of (I-a) with a reagent R^{5a} -W (III)

wherein W is a reactive leaving group and R^{5a} is as R^5 provided that it is not hydrogen, thus obtaining a compound of formula




b) reacting a compound of formula




wherein W is a reactive leaving group, with a cyanide, if desired in the presence of a reaction-inert solvent ;

c) arylating a 5,6-dihydro-1,2,4-triazine-3,5(2H,4H)-dione of formula



with a benzene derivative of formula




wherein W is a reactive leaving group, preferably in a reaction-inert solvent ;

and optionally converting the compounds of formula (I) into each other by a functional group transformation reaction ; and, if desired, converting the compounds of formula (I) into a therapeutically active non-toxic acid-addition, metal or amine substitution salt form by treatment with an appropriate acid or base, or conversely, converting the acid-addition, metal or amine substitution salt into the free base form with alkali or acid ; and/or preparing stereochemically isomeric forms thereof.

18. A process for preparing a chemical compound as claimed in any one of claims 5 to 8, characterized by


a) eliminating a group E from an intermediate triazadione of formula



wherein E represents an electron attracting group, said elimination being carried out at higher temperat-


ures, if desired, in the presence of an acid, said acid being optionally used as a solvent and optionally, if further desired, in the presence of a reaction-inert solvent ;

b) reacting an intermediate of formula



wherein W represents a reactive leaving group, with a cyanide, if desired, in the presence of a reaction-inert solvent ; and optionally, converting the compounds of formula (II-j) into each other by a functional group transformation reaction ; and, if desired, converting the compounds of formula (II-j) into a therapeutically active non-toxic acid-addition, metal or amine substitution salt form by treatment with an appropriate acid or base, or conversely, converting the acid-addition, metal or amine substitution salt into the free base form with alkali or acid ; and/or preparing stereochemically isomeric forms thereof.


19. A process for preparing a chemical compound as claimed in any one of claims 9 to 11, characterised by, adding an organic or inorganic base to an intermediate of formula




or a stereochemically isomeric form thereof.

35 Ansprüche

1. Chemische Verbindung mit der Formel



ein pharmazeutisch annehmbares Säureadditions-, Metall- oder Aminsubstitutionssalz oder eine stereochemisch isomere Form hiervon, worin :
Ar Thienyl, mit Halogen substituiertes Thienyl, Naphthalinyl oder ein Rest der Formel



5

10 ist ;

R für Wasserstoff, C₁₋₆ Alkyl, CycloC₃₋₆ alkyl, Aryl oder (Aryl)C₁₋₆ alkyl steht ;

R¹ Cyano oder einen Rest der Formel -C(=X)-Y-R⁹ darstellt ; welches X für O oder S steht, Y O, S, NR¹⁰ oder eine direkte Bindung darstellt ; R⁹ Wasserstoff, Aryl, C₃₋₆Cycloalkyl oder C₁₋₆ Alkyl ist, gegebenenfalls substituiert mit Aryl, Hydroxy, Amino, Mono- und Di(C₁₋₆ alkyl) amino, Piperidiny, Pyrrolidiny, 4-Morpholiny, Piperaziny, 4-(C₁₋₆ Alkyl)-piperaziny, 4-(C₁₋₆ Alkylcarbonyl)-piperaziny oder 4-((Aryl)C₁₋₆ alkyl)-piperaziny ;

15

und wenn Y eine direkte Bindung darstellt, R⁹ auch Halogen sein kann ; R¹⁰ Wasserstoff, C₁₋₆ Alkyl oder (Aryl)C₁₋₆ alkyl ist ; oder R⁹ und R¹⁰ zusammengekommen mit dem Stickstoffatom, welches die genannten Reste R⁹ und R¹⁰ trägt, einen Piperidiny-, Pyrrolidiny-, 4-Morpholiny-, Piperaziny-, 4-(C₁₋₆ Alkyl)piperaziny-, 4-(C₁₋₆ Alkylcarbonyl)-piperaziny-, 4-(C₁₋₆ Alkylloxycarbonyl)piperaziny- oder einem 4-((Aryl)-C₁₋₆ alkyl)-piperazinyrest bilden können ;

20

R², R³, R⁶, R⁷ und R⁸ jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluomethyl, C₁₋₆ Alkyl, Hydroxy, C₁₋₆ Alkyl-oxy, C₁₋₆ Alkylcarbonyloxy, Mercapto, C₁₋₆ Alkylthio, C₁₋₆ Alkylsulfinyl, (Trifluormethyl)-sulfonyl, Cyano, Nitro, Amino, Mono- und Di(C₁₋₆ alkyl)amino oder (C₁₋₆ Alkylcarbonyl)amino sind ;

25

R⁴ und R⁵ jeweils unabhängig voneinander Wasserstoff, Aryl, CycloC₃₋₆ alkyl, C₂₋₆ Alkenyl, C₂₋₆ Alkiny, (Aryl)C₂₋₆ alkenyl oder C₁₋₆ Alkyl, gegebenenfalls substituiert mit Aryl, Hydroxy, Amino, Mono- und Di(C₁₋₆ Alkyl)amino, Piperidiny, Pyrrolidiny, 4-Morpholiny, Piperaziny, 4-(C₁₋₆ Alkyl)-piperaziny, 4-(C₁₋₆ Alkylcarbonyl)-piperaziny, 4-(C₁₋₆ Alkyl-oxy-carbonyl)-piperaziny oder 4-((Aryl)C₁₋₆ alkyl)piperaziny bedeuten ; und R⁵ auch C₁₋₆ Alkylcarbonyl, C₁₋₆ Alkylloxycarbonyl, (Aryl)C₁₋₆ alkylloxycarbonyl oder (Aryl)carbonyl sein kann ;

30

worin Aryl für Phenyl steht, welches gegebenenfalls mit bis zu 3 Substituenten substituiert ist, welche jeweils unabhängig voneinander von der Gruppe bestehend aus Halogen, C₁₋₆ Alkyl, C₁₋₆ Alkyl-oxy, Trifluormethyl, Hydroxy, Mercapto, C₁₋₆ Alkylthio, C₁₋₆ Alkylsulfonyl, C₁₋₆ Alkylsulfinyl, Trifluormethylsulfonyl, Cyano, Nitro, Amino, Mono- und Di(C₁₋₆ alkyl)amino und C₁₋₆ Alkylcarbonyl)amino ausgewählt sind.

35

2. Chemische Verbindung nach Anspruch 1, worin Ar Halogenthienyl oder einen Rest der Formel (a) bedeutet, worin R⁶ und R⁷ jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluormethyl, C₁₋₆ Alkyl-oxy, Hydroxy oder C₁₋₆ Alkyl sind ; R⁸ Wasserstoff darstellt ; R für Wasserstoff, C₁₋₆ Alkyl, Phenyl oder Halogenphenyl steht ; R² und R³ jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluormethyl oder C₁₋₆ Alkyl sind ; und R⁴ Wasserstoff oder C₁₋₆ Alkyl bedeutet.


40

3. Chemische Verbindung nach Anspruch 2, worin Ar ein Rest der Formel (a) ist, worin R⁶ Halogen bedeutet, R⁷ und R⁸ Wasserstoff sind, R für Wasserstoff oder C₁₋₆ Alkyl steht, R² und R³ unabhängig voneinander Halogen oder Wasserstoff darstellen.

4. Chemische Verbindung nach Anspruch 3, worin R⁶ 4-Chlor R ist, für Wasserstoff steht, R² 2-Chlor bedeutet, R³ 6-Chlor oder Wasserstoff darstellt und R⁴ Wasserstoff ist.

45

5. Chemische Verbindung mit der Formel



50


55

ein pharmazeutisch annehmbares Säureadditions-, Metall- oder Amins substitutionssalz oder eine stereochemisch isomere Form hiervon, worin :

Ar Thienyl, mit Halogen substituiertes Thienyl, Naphtaliny oder ein Rest der Formel



- ist ;
 r für Wasserstoff, C₁₋₆ Alkyl, cycloC₃₋₆ alkyl, Aryl oder (Aryl)C₁₋₆ alkyl steht ;
 r¹ Cyano oder einen Rest der Formel -C(=X)-Y-R⁹ darstellt ; welches X für O oder S steht,
 Y O, S, NR¹⁰ oder eine direkte Bindung darstellt ; R⁹ Wasserstoff, Aryl, C₃₋₆Cycloalkyl oder C₁₋₆ Alkyl ist,
 gegebenenfalls substituiert mit Aryl, Hydroxy, Amino, Mono- und Di (C₁₋₆ alkyl) amino, Piperidinyl, Pyrrolidinyl,
 4-Morpholinyl, Piperazinyl, 4-(C₁₋₆ Alkyl)-piperazinyl, 4-(C₁₋₆ Alkylcarbonyl)-piperazinyl, 4-(C₁₋₆ Alkyl-
 15 oxycarbonyl) oder 4-((Aryl)C₁₋₆ alkyl)-piperazinyl ; und wenn Y eine direkte Bindung darstellt, R⁹ auch Halogen sein kann ; R¹⁰ Wasserstoff, C₁₋₆ Alkyl oder (Aryl)C₁₋₆ alkyl ist ; oder R⁹ und R¹⁰ zusammenge-
 nommen mit dem Stickstoffatom, welches die genannten Reste R⁹ und R¹⁰ trägt, einen Piperidinyl-, Pyrrolidinyl-,
 4-Morpholinyl-, Piperazinyl-, 4-(C₁₋₆ Alkyl)piperazinyl-, 4-(C₁₋₆ Alkylcarbonyl)-piperazinyl-, 4-(C₁₋₆
 20 Alkylloxycarbonyl)piperazinyl- oder einen 4-((Aryl)-C₁₋₆ alkyl)-piperazinylrest bilden können ;
 r², r³, r⁶, r⁷ und r⁸ jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluomethyl, C₁₋₆ Alkyl, Hydroxy,
 C₁₋₆ Alkyl-oxo, C₁₋₆ Alkylcarbonyloxy, Mercapto, C₁₋₆ Alkylthio, C₁₋₆ Alkylsulfonyl, C₁₋₆ Alkylsulfinyl, (Trifluor-
 methyl)-sulfonyl, Cyano, Nitro, Amino, Mono- und Di(C₁₋₆ alkyl)amino oder (C₁₋₆ Alkylcarbonyl)amino sind ;
 r⁴ Wasserstoff, Aryl, CycloC₃₋₆ alkyl, C₂₋₆ Alkenyl, C₂₋₆ Alkynyl, (Aryl)-C₂₋₆ alkenyl oder C₁₋₆ Alkyl gegebenen-
 25 falls substituiert mit Aryl, Hydroxy, Amino, Mono- und Di(C₁₋₆ Alkyl)amino, Piperidinyl, Pyrrolidinyl, 4-Mor-
 pholinyl, Piperazinyl, 4-(C₁₋₆ Alkyl)-piperazinyl, 4-(C₁₋₆ Alkyl)-piperazinyl, 4-(C₁₋₆ Alkylcarbonyl)-piperazinyl,
 4-(C₁₋₆ Alkyl-oxo-carbonyl)-piperazinyl oder 4-((Aryl)C₁₋₆ alkyl)piperazinyl bedeutet ;
 worin Aryl für Phenyl steht, welches gegebenenfalls mit bis zu 3 Substituenten substituiert ist, welche jeweils
 unabhängig voneinander von der Gruppe bestehend aus Halogen, C₁₋₆ Alkyl, C₁₋₆ Alkyl-oxo, Trifluomethyl,
 30 Hydroxy, Mercapto, C₁₋₆ Alkylthio, C₁₋₆ Alkylsulfonyl, C₁₋₆ Alkylsulfinyl, Trifluormethylsulfonyl, Cyano, Nitro,
 Amino, Mono- und Di(C₁₋₆ alkyl)amino und (C₁₋₆ Alkylcarbonyl)amino ausgewählt sind ; mit der Maßgabe,
 daß, wenn r¹ für Cyano steht, eine der folgenden Bedingungen erfüllt wird :
- a) r besitzt eine andere Bedeutung als Wasserstoff, C₁₋₆ Alkyl, CycloC₃₋₆ alkyl oder Aryl ; oder
 - 35 b) Ar' besitzt eine andere Bedeutung als die eines Restes der Formel (a'), worin im genannten Rest der Formel (a') r⁶ eine andere Bedeutung als Wasserstoff, Halogen, Trifluomethyl, C₁₋₆ Alkyl, C₁₋₆ Alkyl-oxo, C₁₋₆ Alkylthio oder C₁₋₆ Alkylsulfonyl aufweist ; oder
 - c) r² besitzt eine andere Bedeutung als Wasserstoff, Halogen, Trifluomethyl oder C₁₋₆ Alkyl ; oder
 - d) r⁴ besitzt eine andere Bedeutung als Wasserstoff.
 - 40 6. Chemische Verbindung nach Anspruch 5, worin Ar' Halogenthienyl oder einen Rest der Formel (a') bedeutet, worin r⁶ und r⁷ jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluomethyl, C₁₋₆ Alkyl-oxo, Hydroxy oder C₁₋₆ Alkyl sind ; r⁸ Wasserstoff darstellt ; r für Wasserstoff, C₁₋₆ Alkyl, Phenyl oder Halogenphenyl steht ; r² und r³ jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluomethyl oder C₁₋₆ Alkyl sind ; und r⁴ Wasserstoff oder C₁₋₆ Alkyl bedeutet.
 - 45 7. Chemische Verbindung nach Anspruch 6, worin Ar' ein Rest der Formel (a') ist, worin r⁶ Halogen bedeutet, r⁷ und r⁸ Wasserstoff sind, r für Wasserstoff oder C₁₋₆ Alkyl steht, r² und r³ unabhängig voneinander Halogen oder Wasserstoff darstellen.
 8. Chemische Verbindung nach Anspruch 7, worin r⁶ 4-Chlor ist, r für Wasserstoff steht, r² 2-chlor bedeutet, r³ 6-Chlor oder Wasserstoff darstellt und r⁴ Wasserstoff ist.
 - 50 9. Pharmazeutisch annehmbares Metall- oder Amins substitutionssalz einer chemischen Verbindung mit der Formel



(II-k),

oder eine stereochemisch isomere Form hiervon, worin R^{6-a} , R^{7-a} und R^{8-a} jeweils unabhängig voneinander Wasserstoff, Halogen, Trifluomethyl, C_{1-6} Alkyl, C_{1-6} Alkyloxy, C_{1-6} Alkylthio oder C_{1-6} Alkylsulfonyl bedeuten; R^{2-a} und R^{3-a} jeweils unabhängig voneinander Wasserstoff, Halogen, trifluomethyl oder C_{1-6} Alkyl darstellen; und R^a Wasserstoff, C_{1-6} Alkyl, Cyclo C_{3-6} Alkyl oder Phenyl ist, das gegebenenfalls mit zu 3 Substituenten substituiert ist, welche unabhängig voneinander aus der Gruppe, bestehend aus Halogen, Trifluomethyl, C_{1-6} Alkyl, C_{1-6} Alkyloxy, C_{1-6} Alkylthio und C_{1-6} Alkylsulfonyl ausgewählt sind.

10. Pharmazeutisch annehmbares Metall- oder Amins substitutionssalz nach Anspruch 9, worin R^{6-a} Halogen bedeutet, R^{7-a} und R^{8-a} Wasserstoff darstellen, R^a Wasserstoff oder C_{1-6} Alkyl ist und R^{2-a} und R^{3-a} unabhängig voneinander Wasserstoff, Halogen, C_{1-6} Alkyl oder C_{1-6} Alkyloxy bedeuten.

11. Pharmazeutisch annehmbares Metall- oder Amins substitutionssalz nach Anspruch 10, worin R^{6-a} für Chlor steht, R^a Wasserstoff bedeutet, R^{2-a} Chlor ist und R^{3-a} Wasserstoff oder Chlor darstellt.

12. Pharmazeutische Zusammensetzung, umfassend einen geeigneten pharmazeutischen Träger und als wirksamen Bestandteil eine therapeutisch wirksame Menge einer Verbindung nach einem der Ansprüche 1 bis 11.

13. Anti-Protozoenzusammensetzung, umfassend einen inerten Träger und als wirksamen Bestandteil eine wirksame anti-Protozoenmenge einer Verbindung nach einem der Ansprüche 1 bis 11.


14. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, dadurch gekennzeichnet, daß eine therapeutisch wirksame Menge einer Verbindung nach einem der Ansprüche 1 bis 11 mit geeigneten pharmazeutischen Trägern innig vermischt wird.

15. Verbindung nach einem der Ansprüche 1 bis 11 zur Verwendung als Arzneimittel.

16. Verbindung nach einem der Ansprüche 1 bis 11 zur Verwendung als Anti-Protozoenarzneimittel.


17. Verbindung zur Herstellung einer chemischen Verbindung nach einem der Ansprüche 1 bis 4, gekennzeichnet durch

a) Reduzieren einer Zwischenverbindung der Formel




(II),

wodurch eine Verbindung der Formel



(I-a) erhalten wird,


gegebenenfalls gefolgt von einer N-Alkylierungsreaktion von (I-a) mit einem Reagens R^{5-a} -W (III), worin W eine reaktive Leavinggruppe ist und R^{5-a} die gleiche Bedeutung wie R^5 besitzt, vorausgesetzt, daß es nicht für Wasserstoff steht, wodurch eine Verbindung der Formel



erhalten wird;


10 erhalten wird ;

b) Umsetzen einer Verbindung der Formel




20 worin W eine reaktive Leavinggruppe ist, mit einem cyanid, wenn gewünscht in Gegenwart eines reaktionsinerten Lösungsmittels ;

c) Arylieren von einem 5,6-Dihydro-1,2,4-triazin-3,5(2H,4H)-dion der Formel




30 mit einem Benzolderivat der Formel



40 worin W eine reaktive Leavinggruppe ist, vorzugsweise in einem reaktionsinerten Lösungsmittel ;
und gegebenenfalls Überführen der Verbindungen der Formel (I) ineinander durch eine Überführungsreaktion für funktionelle Gruppen ; und wenn gewünscht, Überführen der Verbindungen der Formel (I) in eine
therapeutisch wirksame nicht-toxische Säureadditions-, Metall- oder Aminsäureadditions- oder Aminsäureadditions-
45 Behandlung mit einer geeigneten Säure oder Base oder umgekehrt, Überführen des Säureadditions-, Metall- oder Aminsäureadditions-
Metall- oder Aminsäureadditions- in die freie Basenform mit Alkali oder Säure ; und/oder Herstellen stereochemisch isomerer Formen hiervon.


18. Verfahren zur Herstellung einer chemischen Verbindung nach einem der Ansprüche 5 bis 8, gekennzeichnet durch :

55 a) Eliminieren einer Gruppe E aus einer Triazadionzwischenverbindung der Formel




worin E eine elektronenanziehende Gruppe darstellt, welche Eliminierung bei höheren Temperaturen, wenn gewünscht in Gegenwart einer Säure, welche Säure gegebenenfalls als ein Lösungsmittel verwendet wird, und gegebenenfalls, wenn ferner gewünscht, in Gegenwart eines reaktionsinerten Lösungsmittels ausgeführt wird ;

b) Umetzen einer Zwischenverbindung der Formel



worin W eine reaktive Leavinggruppe darstellt, mit einem Cyanid, wenn gewünscht in Gegenwart eines reaktionsinerten Lösungsmittels ; und gegebenenfalls Überführen der Verbindung der Formel (II-j) inienander durch eine Überführungsreaktion für funktionelle Gruppen ; und wenn gewünscht, Überführen der Verbindungen der Formel (II-j) in eine therapeutisch wirksame nicht-toxische Säureaddition-, Metall- oder Aminsäureadditionssalzform durch Behandlung mit einer geeigneten Säure oder Base, oder umgekehrt, Überführen des Säureadditions-, Metall- oder Aminsäureadditionssalzes in die freie Basenform mit Alkali oder Säure ; und/oder Herstellen stereochemisch isomerer Formen hiervon.


19. Verfahren zur Herstellung einer chemischen Verbindung nach einer der Ansprüche 9 bis 11, gekennzeichnet durch Zusetzen einer organischen oder anorganischen Base zu einer Zwischenverbindung der Formel



oder einer stereochemisch isomeren Form hiervon.


Revendications

1. Composé chimique de formule



un de ses sels d'addition d'acides, sels de substitution de métal ou d'amine pharmaceutiquement acceptable ou une de leurs formes stéréochimiquement isomériques où :

Ar est un thiényle, thiényle halo-substitué, naphthalényle ou un radical de formule



R est un hydrogène, alcoyle en C₁₋₆, cycloalcoyle en C₃₋₆, aryle ou (aryl)alcoyle en C₁₋₆ ; R¹ est un cyano ou un radical de formule -C(=X)-Y-R⁹ ;

ledit X représentant O ou S, Y représentant O, S, NR¹⁰ ou une liaison directe ; R⁹ étant un hydrogène, aryle, cycloalcoyle en C₃₋₆ ou alcoyle en C₁₋₆ éventuellement substitué par un aryle, hydroxy, amino, mono- et di(alcoyle en C₁₋₆)amino, pipéridinyle, pyrrolidinyle, 4-morpholinyle, pipérazinyle, 4-(alcoyle en C₁₋₆)-pipérazinyle, 4-(alcoyle en C₁₋₆-carbonyl)-pipérazinyle, 4-(alcoyle en C₁₋₆-carbonyl)-pipérazinyle ou 4-((aryl)alcoyle en C₁₋₆)-pipérazinyle ; et où Y est une liaison directe, R⁹ peut être également un halo ; R¹⁰ est un hydrogène, alcoyle en C₁₋₆ ou (aryl)alcoyle en C₁₋₆ ; ou R⁹ et R¹⁰ ensemble avec l'atome d'azote portant lesdits R⁹ et R¹⁰ peuvent former un pipéridinyle, pyrrolidinyle, 4-morpholinyle, pipérazinyle, 4-(alcoyle en C₁₋₆) pipérazinyle, 4-(alcoyle en C₁₋₆-carbonyl)-pipérazinyle, 4-(alcoyloxy en C₁₋₆-carbonyl)-pipérazinyle ou un 4-((aryl)alcoyle en C₁₋₆)-pipérazinyle ;

R², R³, R⁶, R⁷ et R⁸ représentent chacun indépendamment un hydrogène, halo, trifluorométhyle, alcoyle en C₁₋₆, hydroxy, alcoyloxy en C₁₋₆, alcoyle en C₁₋₆-carbonyloxy, mercapto, alcoylthio en C₁₋₆, alcoyle en C₁₋₆-sulfonyl, alcoyle en C₁₋₆-sulfonyle, (trifluorométhyl)-sulfonyl, cyano, nitro, amino, mono- et di(alcoyle en C₁₋₆)amino, ou (alcoyle en C₁₋₆-carbonyl) amino,

R⁴ et R⁵ représentent chacun indépendamment un hydrogène, aryle, cyclo-alcoyle en C₃₋₆, alcényle en C₂₋₆, alcynyle en C₂₋₆, (aryl)-alcényle en C₂₋₆ ou alcoyle en C₁₋₆ éventuellement substitué par un aryle, hydroxy, amino, mono- et di(alcoyle en C₁₋₆) amino, pipéridinyle, pyrrolidinyle, 4-morpholinyle, pipérazinyle, 4-(alcoyle en C₁₋₆)-pipérazinyle, 4-(alcoyle en C₁₋₆-carbonyl)-pipérazinyle, 4-(alcoyloxy en C₁₋₆-carbonyl)-pipérazinyle, ou 4-((aryl)alcoyle en C₁₋₆)-pipérazinyle ; et R⁵ peut également être un alcoyle en C₁₋₆-carbonyl, alcoyloxy en C₁₋₆-carbonyl, (aryl)-alcoyloxy en C₁₋₆-carbonyl ou (aryl)carbonyl ;


où aryle est un phényle, éventuellement substitué par jusqu'à 3 substituants chacun choisi indépendamment dans le groupe constitué par halo, alcoyle en C₁₋₆, alcoyloxy en C₁₋₆, trifluorométhyle, hydroxy, mercapto, alcoylthio en C₁₋₆, alcoyle en C₁₋₆-sulfonyl, alcoyle en C₁₋₆-sulfonyle, trifluorométhylsulfonyl, cyano, nitro, amino, mono- et di(alcoyle en C₁₋₆)amino et (alcoyle en C₁₋₆-carbonyl)-amino.

2. Composé chimique selon la revendication 1, où Ar est un halothiényle ou un radical de formule (a), où R⁶ et R⁷ représentent chacun indépendamment un hydrogène, halo, trifluorométhyle, alcoyloxy en C₁₋₆, hydroxy ou alcoyle en C₁₋₆ ; R⁸ est un hydrogène ; R est un hydrogène, alcoyle en C₁₋₆, phényle ou halo-phényle ; R² et R³ représentent chacun indépendamment un hydrogène, halo, trifluorométhyle ou alcoyle en C₁₋₆ ; et R⁴ est un hydrogène ou un alcoyle en C₁₋₆.

3. Composé chimique selon la revendication 2, où Ar est un radical de formule (a) où R⁶ est un halo, R⁷ et R⁸ sont un hydrogène, R est un hydrogène ou un alcoyle en C₁₋₆, R² et R³ représentent indépendamment un halo ou un hydrogène.


4. Composé chimique selon la revendication 3, où R⁶ est un 4-chloro, R est un hydrogène, R² est un 2-chloro, R³ est un 6-chloro ou un hydrogène et R⁴ est un hydrogène.

5. Composé chimique de formule



un de ses sels d'addition d'acides, sel de substitution de métal ou d'amine pharmaceutiquement acceptable ou une de leurs formes stéréochimiquement isomériques, où :

Ar' est un thiényle, thiényle halo-substitué, naphthalényle, ou un radical de formule



r est un hydrogène, alcoyle en C₁₋₆, cyclo-alcoyle en C₃₋₆, aryle ou (aryl)-alcoyle en C₁₋₆ ;

r¹ est un cyano ou un radical de formule -C(=X)-Y-R⁹ ;

15 ledit X représentant O ou S,

Y représentant O, S, NR¹⁰ ou une liaison directe ; R⁹ représentant un hydrogène, aryle, cycloalcoyle en C₃₋₆ ou alcoyle en C₁₋₆ éventuellement substitué par un aryle, hydroxy, amino, mono- et di(alcoyle en C₁₋₆)amino, pipéridinyle, pyrrolidinyle, 4-morpholinyle, pipérazinyle, 4-(alcoyle en C₁₋₆) pipérazinyle, 4-(alcoyle en C₁₋₆)-pipérazinyle, 4-(alcoyloxy en C₁₋₆-carbonyl)-pipérazinyle, ou 4-((aryl)-alcoyle en C₁₋₆)-pipérazinyle ; et où

20 Y est une liaison directe, R⁹ peut également être un halo ; R¹⁰ est un hydrogène, alcoyle en C₁₋₆ ou (aryl)-alcoyle en C₁₋₆ ; ou R⁹ et R¹⁰ pris ensemble avec l'atome d'azote portant lesdits R⁹ et R¹⁰ peut former un radical pipéridinyle, pyrrolidinyle, 4-morpholinyle, pipérazinyle, 4-(alcoyle en C₁₋₆)-pipérazinyle, 4-(alcoyle en C₁₋₆-carbonyl)-pipérazinyle, 4-(alcoyloxy en C₁₋₆-carbonyl)-pipérazinyle ou 4-((aryl)-alcoyle en C₁₋₆)-pipérazinyle ;

25 r², r³, r⁶, r⁷ et r⁸ représentent chacun indépendamment un hydrogène, halo, trifluorométhyle, alcoyle en C₁₋₆, hydroxy, alcoyloxy en C₁₋₆, alcoyle en C₁₋₆-carbonyloxy, mercapto, alcoylthio en C₁₋₆, alcoyle en C₁₋₆-sulfonyle, alcoyle en C₁₋₆-sulfinyle, (trifluorométhyl)-sulfonyle, cyano, nitro, amino, mono- et di(alcoyle en C₁₋₆)-amino, ou (alcoyle en C₁₋₆-carbonyl)-amino ;

30 r⁴ est un hydrogène, aryle, cyclo-alcoyle en C₃₋₆, alcényle en C₂₋₆, alcynyle en C₂₋₆, (aryl)-alcényle en C₂₋₆ ou alcoyle en C₁₋₆ éventuellement substitué par un aryle, hydroxy, amino, mono- et di(alcoyle en C₁₋₆)-amino, pipéridinyle, pyrrolidinyle, 4-morpholinyle, pipérazinyle, 4-(alcoyle en C₁₋₆), pipérazinyle 4-(alcoyle en C₁₋₆-carbonyl)-pipérazinyle, 4-(alcoylory en C₁₋₆-carbonyl)-pipérazinyle, ou 4-((aryl)-alcoyle en C₁₋₆)-pipérazinyle ;

où aryle est un phényle, éventuellement substitué par jusqu'à 3 substituants chacun choisi indépendamment dans le groupe constitué par halo, alcoyle en C₁₋₆, alcoyloxy en C₁₋₆, trifluorométhyle, hydroxy, mercapto, alcoylthio en C₁₋₆, alcoyle en C₁₋₆-sulfonyle, alcoyle en C₁₋₆-sulfinyle, trifluorométhylsulfonyle, cyano, nitro, amino, mono- et di(alcoyle en C₁₋₆)-amino et alcoyle en C₁₋₆-carbonyl)-amino

sous réserve que lorsque r¹ est un cyano, l'une des conditions suivantes est satisfaite :

a) r est différent d'un hydrogène, alcoyle en C₁₋₆, cyclo-alcoyle en C₃₋₆ ou aryle ; ou

40 b) Ar' est différent d'un radical de formule (a') où dans ledit radical de formule (a'), r⁶ est différent d'un hydrogène, halo, trifluorométhyle, alcoyle en C₁₋₆, alcoyloxy en C₁₋₆, alcoylthio en C₁₋₆, ou alcoyle en C₁₋₆-sulfonyle ; ou

c) r² est différent d'un hydrogène, halo, trifluorométhyle ou alcoyle en C₁₋₆ ;


d) r⁴ est différent d'un hydrogène.

45 6. Composé chimique selon la revendication 5, où Ar' est un halothiényle ou un radical de formule (a') où r⁶ et r⁷ représentent chacun indépendamment un hydrogène, halo, trifluorométhyle, alcoyloxy en C₁₋₆, hydroxy ou alcoyle en C₁₋₆ ; et r⁸ est un hydrogène, r est un hydrogène, alcoyle en C₁₋₆, phényle ou halo-phényle ; r² et r³ représentent chacun indépendamment un hydrogène, halo, trifluorométhyle ou alcoyle en C₁₋₆ ; et r⁴ est un hydrogène ou un alcoyle en C₁₋₆.

50 7. Composé chimique selon la revendication 6, où Ar' est un radical de formule (a'), où r⁶ est un halo, r⁷ et r⁸ représentent un hydrogène, r est un hydrogène ou un alcoyle en C₁₋₆, r² et r³ représentent chacun indépendamment un halo ou un hydrogène.

8. Composé chimique selon la revendication 7 où r⁶ est un 4-chloro, r est un hydrogène, r² est un 2-chloro, r³ est un 6-chloro ou un hydrogène et r⁴ est un hydrogène.

55 9. Sel de substitution de métal ou d'amine pharmaceutiquement acceptable d'un composé chimique de formule



(II-k).

- 10 ou d'une de ses formes stéréochimiquement isomériques, où R^{6-a} , R^{7-a} et R^{8-a} représentent chacun indépendamment un hydrogène, halo, trifluorométhyle, alcoyle en C_{1-6} , alcoyloxy en C_{1-6} , alcoylthio en C_{1-6} ou alcoyle en C_{1-6} -sulfonyl ; R^{2-a} et R^{3-a} représentent chacun indépendamment un hydrogène, halo, trifluorométhyle ou alcoyle en C_{1-6} ; et R^a est un hydrogène, alcoyle en C_{1-6} , cyclo-alcoyle en C_{2-6} ou phényle éventuellement substitué par jusqu'à trois substituants chacun choisi indépendamment dans le groupe constitué par halo, trifluorométhyle, alcoyle en C_{1-6} , alcoyloxy en C_{1-6} , alcoylthio en C_{1-6} et alcoyle en C_{1-6} -sulfonyloxy.

- 15 10. Sel de substitution de métal ou d'amine pharmaceutiquement acceptable selon la revendication 9, où R^{6-a} est un halo, R^{7-a} et R^{8-a} représentent un hydrogène, R^a représente un hydrogène ou un alcoyle en C_{1-6} et R^{2-a} et R^{3-a} représentent chacun indépendamment un hydrogène, halo, alcoyle en C_{1-6} ou alcoyloxy en C_{1-6} .

- 20 11. Sel de substitution de métal ou d'amine pharmaceutiquement acceptable selon la revendication 10, où R^{6-a} est un chloro, R^a est un hydrogène, R^{2-a} est un chloro, et R^{3-a} est un hydrogène ou un chloro.

12. Composition pharmaceutique comprenant un support pharmaceutiquement acceptable et comme ingrédient actif une quantité thérapeutiquement efficace d'un composé selon l'une quelconque des revendications 1-11.

- 25 13. Composition antiprotozoaire comprenant un support inerte et comme ingrédient actif une quantité antiprotozoaire efficace d'un composé selon l'une quelconque des revendications 1-11.


14. Procédé de préparation d'une composition pharmaceutique, caractérisé en ce qu'on mélange intimement une quantité thérapeutiquement efficace d'un composé selon l'une quelconque des revendications 1-11 avec des supports pharmaceutiques appropriés.

- 30 15. Composé selon l'une quelconque des revendications 1-11 pour application comme médicament.

16. Composé selon l'une quelconque des revendications 1-11 pour application comme médicament antiprotozoaire.


17. Procédé de préparation d'un composé chimique selon l'une quelconque des revendications 1-4, caractérisé en ce que

- 35 a) on réduit un intermédiaire de formule



(II)

- 45 obtenant ainsi un composé de formule




(I-a).

55


ceci étant éventuellement suivi par une réaction de N-alcoylation de (I-a) avec un réactif



5 où W est un groupe sortant réactif, et R^{5-a} est tel que R^5 à condition qu'il ne soit pas un hydrogène donnant ainsi un composé de formule




b) on fait réagir un composé de formule




30 où W est un groupe sortant réactif, avec un cyanure, si on le désire en présence d'un solvant inerte vis-à-vis de la réaction ;

c) on aryle une 5,6-dihydro-1,2,4-triazine-3,5(2H,4H)-dione de formule



avec un dérivé de benzène de formule




50 où W est un groupe sortant réactif, de préférence dans un solvant inerte vis-à-vis de la réaction ; et le cas échéant, on transforme les composés de formule (I) les uns dans les autres par une réaction de transformation de groupes fonctionnels ; et, si on le désire, on transforme les composés de formule (I) en un sel d'addition d'acides, ou sel de substitution de métal ou d'amine non toxique thérapeutiquement actif par traitement avec un acide ou une base appropriée, ou inversement, on transforme le sel d'addition d'acides, sel de substitution de métal ou d'amine sous la forme base libre avec une base ou un acide ; et/ou par leurs formes stéréochimiquement isomériques.

18. procédé de préparation d'un composé chimique selon l'une quelconque des revendications 5 à 8,


caractérisé en ce que

a) on élimine un groupe E d'une triazadione intermédiaire de formule




où E représente un groupe attracteur d'électrons, ladite élimination étant conduite à des températures plus élevées, si on le désire, en présence d'un acide, ledit acide étant éventuellement utilisé comme solvant et, le cas échéant, si on le désire, en outre, en présence d'un solvant inerte vis-à-vis de la réaction ;

b) on fait réagir un intermédiaire de formule



où W représente un groupe sortant réactif, avec un cyanure, si on le désire, en présence d'un solvant inerte vis-à-vis de la réaction ; et le cas échéant, on transforme les composés de formule (II-j) les uns dans les autres par une réaction de transformation de groupes fonctionnels ; et, si on le désire, on transforme les composés de formule (II-j) en une forme sel d'addition d'acides, de substitution de métal ou d'amine non toxique thérapeutiquement active, par traitement avec un acide ou une base appropriée, ou, inversement, on transforme le sel d'addition d'acides, de substitution de métal ou d'amine en la forme base libre avec une base ou un acide ; et/ou on prépare leurs formes stéréochimiquement isomériques.

19. procédé de préparation d'un composé chimique selon l'une quelconque des revendications 9-11, caractérisé en ce qu'on ajoute une base organique ou inorganique à un intermédiaire de formule



ou une de ses formes stéréochimiquement isomériques.